

Express Mail™ Label No. EL921061407US

Date of Deposit: December 19, 2001

I hereby certify that this is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above, addressed to: Assistant Commissioner for Patents Washington, Box Patent Application, D.C. 20231

By: Tracie Brooks

Tracie Brooks

Docket No: **407T-898010US**

Client Ref: **2000-072-1**

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

U.S. Patent Application For

A NOVEL GENE ASSOCIATED WITH REGULATION OF
ADIPOSY AND INSULIN RESPONSE

Inventor(s): **KAREN REUE**, a citizen of the United States of America, residing at 20338 Eastwood Ave. Torrance CA 90503

MIKLÓS PÉTERFY, a citizen of Hungary, residing at 637 Levering Ave., #3, Los Angeles, CA 90024

Assignee: **The Regents of the University of California**

Entity: **Small Entity**

LAW OFFICES OF JONATHAN ALAN QUINE

2033 Clement Ave

Alameda, CA 94501

Tel: (510) 337-7871

Fax: (510) 337-7877

A NOVEL GENE ASSOCIATED WITH REGULATION OF ADIPOSIITY AND INSULIN RESPONSE

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to and benefit of USSN 60/257,772, filed
5 on 22 December 2000, which is incorporated herein by reference in its entirety for all
purposes.

STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

[0002] This invention was made with Government support by the Veterans
10 Administration, and under Grant No: HL28481, awarded by the National Institutes of
Health. The Government of the United States of America may have certain rights in this
invention.

FIELD OF THE INVENTION

[0003] This invention pertains to the field of adipose tissue development and fat
15 metabolism. In particular, this invention pertains to the identification of a gene implicated
in the development of lipodystrophy.

BACKGROUND OF THE INVENTION

[0004] Considerable work has been performed in the field of adipocyte
differentiation in an effort to delineate the specific factors and processes involved. Fat
20 cells or adipocytes are a specialized cell type that synthesizes and stores fat (triglycerides)
in periods of nutritional abundance, and hydrolyze these fats when needed to meet demand
for energy. The size and distribution of adipose tissue stores in humans and animals
clearly influences metabolism and the development of diseases including, but not limited
to obesity and diabetes.

25 [0005] The development of mature adipocytes from precursor cells, a process
known as differentiation has been widely studied. Most adipocyte differentiation occurs

shortly before or after birth, but further differentiation can occur at any time during life in response to various hormonal and nutritional signals.

[0006] A few key proteins that are expressed early in the process of differentiation and are required for adipocyte development have been identified. These include the nuclear hormone receptor peroxisome proliferator-activated receptor γ (PPAR γ) and CAAT/enhancer binding protein α (C/EBP α). It is known that these proteins function as transcription activators to induce the expression of genes required for conversion of precursor cells into mature adipocytes. However, many aspects of the regulation and function of these proteins have not been delineated. It is known that proteins such as PPAR γ and C/EBP α function in complexes with other protein factors, known as co-activators, to effect transcription.

SUMMARY OF THE INVENTION

[0007] This invention pertains to the identification and isolation of a gene implicated in the fatty liver dystrophy (*fld*) phenotype. In particular, this invention relates to the isolation of both mouse and human forms of a novel gene (designated herein as *Lpin1* and *LPIN1*, respectively), that is rearranged in the *fld* genome leading to a null allele. An *fld* mutant animal exhibits an array of abnormalities in lipid and glucose metabolism. In particular, as *fld* animals age, they exhibit reduced weight gain and adipose tissue mass, develop glucose intolerance and hyperinsulinemia, and are more susceptible to diet-induced atherosclerosis. Without being bound by a particular theory, it is believed that a *Lpin1* gene product may interact with proteins such as PPAR γ and C/EBP α , or other proteins involved in adipocyte differentiation.

[0008] In one embodiment, this invention provides a method of screening for an agent that alters adipose tissue development. The method involves contacting a cell comprising a *Lpin1* gene with a test agent; and detecting a change in the expression or activity of a *Lpin1* gene product as compared to the expression or activity of a *Lpin1* gene product in a cell that is contacted with the test agent at a lower concentration, where a difference in the expression or activity of lipin in the contacted cell and the cell that is contacted with the lower concentration indicates that the test agent alters adipose tissue development. In preferred embodiments, the lower concentration is the absence of the test

agent. In certain embodiments, the of *Lpin1* gene product is detected by detecting *Lpin1* mRNA in the sample, *e.g.*, by hybridizing said mRNA to a probe that specifically hybridizes to a *Lpin1* nucleic acid. Preferred hybridization assays include, but are not limited to Northern blot, Southern blot using DNA derived from the *Lpin1* RNA, a array hybridization, affinity chromatography, and *in situ* hybridization. In one particularly preferred embodiment, the assay utilizes a *Lpin1* probe that is a member of a plurality of probes that forms an array of probes (*e.g.* a high-density array). In certain other nucleic acid-based assays, the *Lpin1* mRNA is measured using a nucleic acid amplification reaction.

[0009] In other embodiments, the amount of *Lpin1* gene product is detected by detecting the level of a lipin protein in the biological sample, *e.g.* via a method selected from the group consisting of capillary electrophoresis, Western blot, mass spectroscopy, ELISA, immunochromatography, immunohistochemistry, and the like. The cells used in these assays can be cells *in vivo* in an organism or can be *ex vivo* (*e.g.* cultured cells). In certain embodiments, the test agent is contacted to an animal comprising a cell containing the *Lpin1* nucleic acid or the lipin protein.

[0010] This invention also provides methods of prescreening for an agent that alters adipose tissue development. Such methods typically involve contacting a *Lpin1* nucleic acid or a lipin protein with a test agent; and detecting specific binding of the test agent to the lipin protein or nucleic acid. The methods can further involve recording test agents that specifically bind to the *Lpin1* nucleic acid or protein in a database of candidate agents that alter adipose tissue development. In certain embodiments, the test agent is not an antibody and/or not a protein, and/or not a nucleic acid. Particularly preferred test agents include, but are not limited to small organic molecules. In preferred embodiments, the detecting comprises detecting specific binding of the test agent to the *Lpin1* nucleic acid (*e.g.* via Northern blot, Southern blot using DNA derived from a *Lpin1* RNA, array hybridization, affinity chromatography, *in situ* hybridization, and the like). In other embodiments, the detecting comprises detecting specific binding of the test agent to lipin protein (*e.g.* via capillary electrophoresis, Western blot, mass spectroscopy, ELISA, immunochromatography, immunohistochemistry, and the like). In certain embodiments, the test agent is contacted directly to the *Lpin1* nucleic acid or to the lipin protein, while in other embodiments, the test agent is contacted to a cell containing (comprising) the *Lpin1*

nucleic acid or the lipin protein. The cells can be *in vivo* or *ex vivo*, but particularly preferred cells are *ex vivo* (e.g. cultured fresh cells or cell lines). In certain embodiments, the test agent is contacted to an animal comprising a cell containing (comprising) the *Lpin1* nucleic acid or the lipin protein.

- 5 [0011] This invention also provides an isolated nucleic acid comprising a nucleic acid selected from the group consisting of: a nucleic acid encoding a polypeptide selected from the group consisting of human lipin 1A (SEQ ID NO:3), mouse lipin 1A (SEQ ID NO:4), and mouse lipin 1B (SEQ ID NO:5); a nucleic acid that hybridizes to a nucleic acid selected from the group consisting of mouse *Lpin1* (SEQ ID NO:1), and human
- 10 *LPIN1* (SEQ ID NO2) under stringent conditions; a nucleic acid having the sequence of a nucleic acid selected from the group consisting of mouse *Lpin1* (SEQ ID NO:1), and human *LPIN1* (SEQ ID NO2); a nucleic acid that hybridizes to a nucleic acid selected from the group consisting of mouse *Lpin1* (SEQ ID NO:1), and human *LPIN1* (SEQ ID NO2) under stringent conditions and that encodes an lipin polypeptide; a nucleic acid
- 15 having the sequence of a nucleic acid amplified using primer 1 (SEQ ID NO:6) and primer 2 (SEQ ID NO:7) using cDNA from mouse cells or tissues as a template; and a nucleic acid having the sequence of a nucleic acid amplified using primer 3 (SEQ ID NO:8) and primer 4 (SEQ ID NO:9) using cDNA from human cells or tissues as a template.
- Preferred nucleic acids are sufficiently long to hybridize specifically with a *Lpin1* nucleic acid. A preferred nucleic acid is at least 15 nucleotides in length. In one particularly
- 20 preferred embodiment the nucleic acid comprises a mouse *Lpin1* (SEQ ID NO:1), and/or a human *Lpin1* (SEQ ID NO2).

- [0012] This invention also provides a comprising a polypeptide encoded by a nucleic acid as described above and herein and/or an antibody that specifically binds such
- 25 a polypeptide. Particularly preferred polypeptides include lipin polypeptides comprising a polypeptide that comprises an NLIP domain and a CLIP domain.

- [0013] In still another embodiment, this invention provides a transgenic animal comprising a recombinantly modified *Lpin1/LPIN1* gene such that the recombinantly modified gene does not transcribe a functional lipin protein. A preferred transgenic
- 30 animal is homozygous for the recombinantly modified *Lpin1/LPIN1* gene. Certain preferred transgenic animals are chimeric for cells comprising said recombinantly

modified *Lpin1/LPIN1* gene. Preferred transgenic animals include, felines, canines, equines, largomorphs, murines, mice, rodents, and the like.

[0014] This invention provides a method of identifying a predilection to developing one or more symptoms of lipodystrophy, obesity, diabetes, or atherosclerosis (e.g., a methods of identification mutations in the *LPIN1* gene). Preferred methods involve obtaining a biological sample from the organism; and detecting a mutation in a *Lpin1/LPIN1* gene or gene product from the biological sample. In preferred embodiments, the mutation is an insertion, a deletion, a missense point mutation, a nonsense point mutation, *etc.* Preferred detection methods include, but are not limited to Southern blot, DNA amplification, comparative genomic hybridization (CGH), immunohistochemistry, cytogenetics, and the like. In other embodiments, the detecting comprises detecting a mutation in a polypeptide, e.g. via capillary electrophoresis, Western blot, mass spectroscopy, ELISA, immunochromatography, immunohistochemistry, and the like.

[0015] In still another embodiment, this invention provides a method of identifying a predilection to developing one or more symptoms of lipodystrophy, obesity, diabetes, by detecting alterations in *lpin* expression. Thus the methods preferably involve obtaining a biological sample from the organism; and detecting a *LPIN1* gene product wherein a difference in the amount or activity of the *LPIN1* gene product from the organism as compared to the *LPIN1* gene product from a normal healthy organism indicates that the organism has or is susceptible to lipodystrophy, obesity, diabetes, atherosclerosis, and related pathologies. In preferred embodiments, the amount of *LPIN1* gene product is detected by detecting *LPIN1* mRNA in the cell (e.g., by hybridizing the mRNA or a nucleic acid derived therefrom to a probe that specifically hybridizes to a *LPIN1* nucleic acid). Preferred hybridization methods include, but are not limited to Northern blot, Southern blot using DNA derived from the *LPIN1* RNA, array hybridization, affinity chromatography, *in situ* hybridization, and the like. In certain embodiments, the probe is a member of a plurality of probes that forms an array of probes. In certain embodiments, the level of *LPIN1* mRNA is measured using a nucleic acid amplification reaction. The amount of *LPIN1* gene product can also be detected by detecting the level of lipin protein in the biological sample (e.g. via Western blot, mass spectroscopy, ELISA, immunochromatography, immunohistochemistry, *etc.*).

[0016] This invention also provides methods of mitigating a symptom of lipodystrophy, obesity, diabetes, atherosclerosis or related pathology. The methods involve modulating the concentration and/or activity of a *LPIN1* gene product in a cell of an organism *e.g.*, by upregulating or repressing expression of a heterologous *LPIN1* nucleic acid or by upregulating or repressing expression of an endogenous *LPIN1* gene. In preferred embodiments, lipin expression is upregulated, *e.g.* by transfecting the cell with a vector (*e.g.* constitutive, inducible, tissue-specific, *etc.*) that expresses a lipin protein. Preferred cells are adipocytes.

Definitions.

[0017] The terms "polypeptide", "peptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical analogue of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers.

[0018] The terms "nucleic acid" or "oligonucleotide" or grammatical equivalents herein refer to at least two nucleotides covalently linked together. A nucleic acid of the present invention is preferably single-stranded or double stranded and will generally contain phosphodiester bonds, although in some cases, as outlined below, nucleic acid analogs are included that may have alternate backbones, comprising, for example,

phosphoramidate (Beaucage *et al.* (1993) *Tetrahedron* 49(10):1925) and references therein; Letsinger (1970) *J. Org. Chem.* 35:3800; Sprinzl *et al.* (1977) *Eur. J. Biochem.* 81: 579; Letsinger *et al.* (1986) *Nucl. Acids Res.* 14: 3487; Sawai *et al.* (1984) *Chem. Lett.* 805, Letsinger *et al.* (1988) *J. Am. Chem. Soc.* 110: 4470; and Pauwels *et al.* (1986) *Chemica Scripta* 26: 141 9), phosphorothioate (Mag *et al.* (1991) *Nucleic Acids Res.* 19:1437; and U.S. Patent No. 5,644,048), phosphorodithioate (Briu *et al.* (1989) *J. Am. Chem. Soc.* 111 :2321, O-methylphosphoroamidite linkages (*see* Eckstein, *Oligonucleotides and Analogues: A Practical Approach*, Oxford University Press), and peptide nucleic acid backbones and linkages (*see* Egholm (1992) *J. Am. Chem. Soc.* 114:1895; Meier *et al.* (1992) *Chem. Int. Ed. Engl.* 31: 1008; Nielsen (1993) *Nature*, 365: 566; Carlsson *et al.* (1996) *Nature* 380: 207). Other analog nucleic acids include those with positive backbones (Denpcy *et al.* (1995) *Proc. Natl. Acad. Sci. USA* 92: 6097; non-ionic

backbones (U.S. Patent Nos. 5,386,023, 5,637,684, 5,602,240, 5,216,141 and 4,469,863; Angew. (1991) *Chem. Intl. Ed. English* 30: 423; Letsinger et al. (1988) *J. Am. Chem. Soc.* 110:4470; Letsinger et al. (1994) *Nucleoside & Nucleotide* 13:1597; Chapters 2 and 3, ASC Symposium Series 580, "Carbohydrate Modifications in Antisense Research", Ed. Y.S. Sanghui and P. Dan Cook; Mesmaeker et al. (1994), *Bioorganic & Medicinal Chem. Lett.* 4: 395; Jeffs et al. (1994) *J. Biomolecular NMR* 34:17; *Tetrahedron Lett.* 37:743 (1996)) and non-ribose backbones, including those described in U.S. Patent Nos. 5,235,033 and 5,034,506, and Chapters 6 and 7, ASC Symposium Series 580, *Carbohydrate Modifications in Antisense Research*, Ed. Y.S. Sanghui and P. Dan Cook.

Nucleic acids containing one or more carbocyclic sugars are also included within the definition of nucleic acids (see Jenkins et al. (1995), *Chem. Soc. Rev.* pp169-176). Several nucleic acid analogs are described in Rawls, C & E News June 2, 1997 page 35. These modifications of the ribose-phosphate backbone may be done to facilitate the addition of additional moieties such as labels, or to increase the stability and half-life of such molecules in physiological environments.

[0019] As used herein, the term "derived from a nucleic acid" (e.g., an mRNA) refers to a nucleic acid or protein nucleic acid for whose synthesis the referenced nucleic acid or a subsequence thereof has ultimately served as a template. Thus, a cDNA reverse transcribed or RT-PCR'd from an mRNA, an RNA transcribed from that cDNA, a DNA amplified from the cDNA, an RNA transcribed from the amplified DNA, etc., are all derived from the mRNA. In preferred embodiments, detection of such derived products is indicative of the presence and/or abundance of the original nucleic acid in a sample.

[0020] A "nucleic acid derived from a *Lpin1* gene or cDNA" refers to a nucleic acid whose synthesis the *Lpin1* gene or cDNA has ultimately served as a template. Thus, for example, a cDNA reverse transcribed from a *Lpin1* mRNA, an RNA transcribed from that cDNA, a DNA amplified from the cDNA, an RNA transcribed from the amplified DNA, etc., are all nucleic acids derived from the *Lpin1* gene or cDNA.

[0021] A *Lpin1* nucleic acid refers to a nucleic acid derived from a *Lpin1* gene, mRNA or cDNA, or a nucleic acid having the same sequence as a nucleic acid derived from a *lpin* gene, mRNA or cDNA. A *lpin* nucleic acid also includes fragments of such nucleic acids. In preferred embodiments, the fragments are of sufficient length to uniquely

identify them as lpin gene mRNA or cDNA subsequences. Preferred fragments are at least 10 nucleotides, more preferably at least 15 nucleotides, still more preferably at last 20 nucleotides, and most preferably at least 25, 50, 100 or 200 nucleotides in length.

[0022] The term "antibody", as used herein, includes various forms of modified or altered antibodies, such as an intact immunoglobulin, an Fv fragment containing only the light and heavy chain variable regions, an Fv fragment linked by a disulfide bond (Brinkmann *et al.* (1993) *Proc. Natl. Acad. Sci. USA*, 90: 547-551), an Fab or (Fab)'2 fragment containing the variable regions and parts of the constant regions, a single-chain antibody and the like (Bird *et al.* (1988) *Science* 242: 424-426; Huston *et al.* (1988) *Proc. Nat. Acad. Sci. USA* 85: 5879-5883). The antibody may be of animal (especially mouse or rat) or human origin or may be chimeric (Morrison *et al.* (1984) *Proc Nat. Acad. Sci. USA* 81: 6851-6855) or humanized (Jones *et al.* (1986) *Nature* 321: 522-525, and published UK patent application #8707252).

[0023] The terms "binding partner", or "capture agent", or a member of a "binding pair" refers to molecules that specifically bind other molecules to form a binding complex such as antibody-antigen, lectin-carbohydrate, nucleic acid-nucleic acid, biotin-avidin, *etc.*

[0024] The term "specifically binds", as used herein, when referring to a biomolecule (e.g., protein, nucleic acid, antibody, etc.), refers to a binding reaction which is determinative of the presence of a biomolecule in a heterogeneous population of molecules (e.g., proteins and other biologics). Thus, under designated conditions (e.g. immunoassay conditions in the case of an antibody or stringent hybridization conditions in the case of a nucleic acid), the specified ligand or antibody binds to its particular "target" molecule and does not bind in a significant amount to other molecules present in the sample.

[0025] The terms "hybridizing specifically to" and "specific hybridization" and "selectively hybridize to," as used herein refer to the binding, duplexing, or hybridizing of a nucleic acid molecule preferentially to a particular nucleotide sequence under stringent conditions. The term "stringent conditions" refers to conditions under which a probe will hybridize preferentially to its target subsequence, and to a lesser extent to, or not at all, to other sequences. Stringent hybridization and stringent hybridization wash conditions in the context of nucleic acid hybridization are sequence dependent, and are different under

different environmental parameters. An extensive guide to the hybridization of nucleic acids is found in, *e.g.*, Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Acid Probes part I, chapt 2, Overview of principles of hybridization and the strategy of nucleic acid probe assays*, Elsevier, NY (Tijssen). Generally, highly stringent hybridization and wash conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Very stringent conditions are selected to be equal to the T_m for a particular probe. An example of stringent hybridization conditions for hybridization of complementary nucleic acids which have more than 100 complementary residues on an array or on a filter in a Southern or northern blot is 42°C using standard hybridization solutions, *e.g.*, containing formamide (*see, e.g.*, Sambrook (1989) *Molecular Cloning: A Laboratory Manual (2nd ed.) Vol. 1-3*, Cold Spring Harbor Laboratory, Cold Spring Harbor Press, NY, and detailed discussion, below), with the hybridization being carried out overnight. An example of highly stringent wash conditions is 0.15 M NaCl at 72°C for about 15 minutes. An example of stringent wash conditions is a 0.2x SSC wash at 65°C for 15 minutes (*see, e.g.*, Sambrook *supra.*) for a description of SSC buffer). Often, a high stringency wash is preceded by a low stringency wash to remove background probe signal. An example medium stringency wash for a duplex of, *e.g.*, more than 100 nucleotides, is 1x SSC at 45°C for 15 minutes. An example of a low stringency wash for a duplex of, *e.g.*, more than 100 nucleotides, is 4x to 6x SSC at 40°C for 15 minutes.

[0026] The term "test agent" refers to an agent that is to be screened in one or more of the assays described herein. The agent can be virtually any chemical compound. It can exist as a single isolated compound or can be a member of a chemical (*e.g.* combinatorial) library. In a particularly preferred embodiment, the test agent will be a small organic molecule.

[0027] The term "small organic molecule" refers to a molecule of a size comparable to those organic molecules generally used in pharmaceuticals. The term excludes biological macromolecules (*e.g.*, proteins, nucleic acids, *etc.*). Preferred small

organic molecules range in size up to about 5000 Da, more preferably up to 2000 Da, and most preferably up to about 1000 Da.

[0028] The term database refers to a means for recording and retrieving information. In preferred embodiments the database also provides means for sorting and/or searching the stored information. The database can comprise any convenient media including, but not limited to, paper systems, card systems, mechanical systems, electronic systems, optical systems, magnetic systems or combinations thereof. Preferred databases include electronic (e.g. computer-based) databases. Computer systems for use in storage and manipulation of databases are well known to those of skill in the art and include, but are not limited to "personal computer systems", mainframe systems, distributed nodes on an inter- or intra-net, data or databases stored in specialized hardware (e.g. in microchips), and the like.

BRIEF DESCRIPTION OF THE DRAWINGS

[0029] Figures 1A, 1B, 1C, and 1D illustrate adipose deficiency in *fld* mutant mice. Figure 1A: Exposed ventral view of wild-type and *fld* littermates at 3 months of age showing vastly reduced epididymal adipose tissue in the *fld* mouse. Figure 1B and Figure 1C: Cross sections of epididymal fat pads from wild-type and *fld* mice, respectively. Tissue was stained with haematoxylin/eosin and viewed at a magnification of 330x. Adipocytes from *fld* mice appear immature, with small heterogeneous lipid droplets. Figure 1D: Plasma leptin levels are substantially reduced in *fld* compared to wild-type (wt) mice. Values represent mean \pm SD for three mice of each type. Asterisk, $P < 0.005$.

[0030] Figures 2A through 2D illustrate the structure of *fld* and *fld*^{2J} mutant alleles. Figure 2A: Genomic DNA from wild-type (+/+), *fld* (*fld/fld*) and heterozygous (+/*fld*) mice was restriction digested with *Hind*III (left panel) or *Sac*I (right panel) and hybridized to probes from the 5' or 3' region of lipin cDNA (see panel c for position of probes in the gene). The *fld* allele exhibits aberrant hybridization patterns indicating DNA deletion and duplication for the 5' and 3' probes, respectively. Figure 2B: Genomic DNA from wild-type (w) and *fld/fld* (f) mice was PCR amplified using primer pairs flanking the regions that are deleted (p1 + p2) and duplicated (p3 + p4) in the *fld* mutant allele. PCR products were generated from the wild-type allele with primer pairs p1 + p2 and p3 + p4, whereas

products from the *fld* allele were only generated when the two forward primers (p1 + p3) or the two reverse primers (p2 + p4) were paired, confirming an internal gene rearrangement. Figure 2C: Structure of wild-type and *fld* alleles of *Lpin1*. At top, exons are represented as boxes, with filled and open boxes representing coding and non-coding regions, respectively. Below is a schematic diagram of the wild-type allele, where the gene is divided into 5 segments represented by distinctly colored rectangles, with arrowheads indicating orientation. In the rearranged *fld* allele shown at bottom, the red segment of the wild-type allele containing exons 2 and 3 has been deleted, while the green segment containing most of the coding exons has been inverted (indicated by reverse arrowhead). Moreover, in *fld* the yellow segment of the wild-type allele containing a portion of the 3'UTR has been both duplicated and inverted. Positions of the probes and primers used in experiments presented in Figures 2A and 2B are also indicated. Figure is not drawn to scale. Figure 2D: Sequence traces showing the Gly84Arg substitution in the *fld^{2J}* allele. The mutation occurs in the NLIP domain of the protein in a position that is invariably conserved in species ranging from mouse and human to yeast (see asterisk in Figure 4D).

[0031] Figures 3A through 3C illustrate lipin mRNA expression. Figure 3A: Lipin mRNA tissue distribution in wild-type mouse tissues. A partial lipin cDNA probe (exons 1-8) was hybridized to 20 μ g total RNA (all tissues except liver), or 2 μ g poly A+ RNA (liver). Figure 3B: Aberrant lipin mRNA expression levels in adipose tissue from *fld/fld* and *fld^{2J}/fld^{2J}* mice. At left, no full length mRNA was detected in *fld* tissue; at right, elevated lipin mRNA was detected in *fld^{2J}* tissue. Ethidium bromide stained RNA is shown below blots to demonstrate equal loading. Figure 3C: Lipin mRNA induction during 3T3-L1 preadipocyte differentiation. Total RNA (20 μ g) from cultures of 3T3-L1 cells maintained in differentiation medium for 0 to 6 days (see Methods) was hybridized to cDNA for lipin or adipsin.

[0032] Figures 4A through 4C illustrate evolutionary conservation of the lipin protein family and nuclear localization of lipin. Figure 4A: Lipin-homologs were identified from mouse, human, *Drosophila melanogaster* (D.m.), *Caenorhabditis elegans* (C.e.), *Saccharomyces cerevisiae* (S.c.), *Schizosaccharomyces pombe* (S.p.), *Arabidopsis thaliana* (A.t.) and *Plasmodium falciparum* (P.f.). Lpin1 (GenBank accession number AAF44296), Lpin2 and Lpin3 sequences were deduced from full-length cDNAs obtained

by RACE cloning in this study. The LPIN1 (Q14693), LPIN2 (Q92539), D.m. (AAF59125), C.e. (CAA16154), S.c. (P32567), S.p. (CAB52577), A.t. (AAF23287) and P.f. (CAB10579) protein sequences are based on predictions from EST and genomic sequences. LPIN3 was predicted in this study using the genomic sequence in AL132654.

5 NLIP (blue) and CLIP (green) domains, and predicted nuclear localization signals (red) are indicated. Figure 4B: 3T3-L1 cells were transfected with pEGFP, a GFP expression vector, or the same vector containing lipin cDNA to produce a lipin-GFP fusion protein. At 48 hr post-transfection, nuclei were stained with the DNA-specific dye Hoechst-33258, and cells were observed in a fluorescent microscope for either GFP (left) or Hoechst

10 fluorescence (right). Magnification is 500 x. Figure 4C: Phylogenetic relationships among the mouse and human lipin protein family members. Multiple sequence alignment and phylogenetic tree were calculated using the ClustalX (Thompson *et al.* (1994) *Nucleic Acids Res.* 22: 4673-4680) and displayed by the Njplot (Perrieren and Gouy (1996) *Biochimie* 78: 364-369) programs. Tree topology is corroborated by 100% bootstrap

15 support (1000 replicates) at all nodes. Homologous chromosomal locations of the corresponding orthologous gene pairs in the mouse (MMU) and human (HSA) genomes are indicated in parentheses. *Lpin1* was initially mapped in the *fld* mutant strain (Rowe *et al.* (1996) *Mamm. Genome* 7: 555-557), and *Lpin2* and *Lpin3* were mapped in this study using a radiation hybrid panel (see Example 1). Genomic localization of the human *LPIN*

20 genes was determined from database resources (reference 26 and the GenBank annotation for AL031667). Figure 4D: Alignments of the NLIP and CLIP domains showing positions of residues that are identical (black background) or similar (gray background) in at least 90% of the sequences. The asterisk above the NLIP sequence denotes the position of the Gly84Arg mutation in *fld^{2J}* mice.

25 [0033] Figures 5A through 5C illustrate that lipin deficiency attenuates obesity in a genetic model of obesity. Figure 5A shows that daily food consumption is the same in ob and fld-ob mice. Figure 5B shows that weight gain is reduced in fld-ob compared to ob mice. Figure 5C illustrates an ob and an fld-ob mouse.

[0034] Figure 6 illustrates the cellular localization of lipin.

30 [0035] Figure 7 illustrates that within the nucleus, lipin occurs within nuclear speckles, a compartment associated with factors involved in nuclear mRNA metabolism.

[0036] Figure 8 illustrates the effect of various lipin mutations on the cellular localization of the mutated lipin.

DETAILED DESCRIPTION

[0037] This invention pertains to the identification and isolation of a gene implicated in the fatty liver dystrophy (fld) phenotype. An *fld* mutant mouse exhibits an array of abnormalities in lipid and glucose metabolism. During the neonatal period (1-10 days), *fld* mice have hypertriglyceridemia and a triglyceride filled, fatty liver. These features disappear by the age of 18-21 days. At this point, other abnormalities appear and persist throughout the animal's lifetime. Most notable is reduced weight gain and failure to accumulate normal amounts of adipose tissue. The *fld* mice also develop glucose intolerance and hyperinsulinemia, a whole body tremor, and have increased susceptibility to atherosclerosis; it is believed that these represent secondary manifestations of the impaired *adipose tissue development and metabolism.

[0038] It is believed that, prior to this invention the genes implicated in the *fld* phenotype were unknown. This invention relates to the isolation of both mouse and human forms of a novel gene, designated herein as *Lpin1* and *LPIN1* for mouse and human, respectively, that is rearranged in the *fld* genome leading to a null allele.

[0039] This gene and/or the mRNA, and/or the protein it encodes provide good targets for screening for agents that modulate adipocyte differentiation. Such agents are expected to be useful in the treatment of obesity, diabetes, and/or related pathologies. In addition, modulators are expected to be effective in regulating fat accumulation and hence weight gain or loss. It is also believed that one or more symptoms of lipodystrophy can be mitigated by increasing levels of an *Lpin1* gene product in an organism.

[0040] Conversely it is believed that, at least in certain cases (*e.g.* pathologies characterized by Overexpression of *lpin*) fat accumulation can be diminished by inhibiting expression or activity of lipin.

I. Assays for agents that modulate *Lpin* expression.

[0041] As indicated above, in one aspect, this invention pertains to the discovery of genes (*e.g.* *Lpin1*) whose inactivation results in a lipodystrophic phenotype. The *Lpin1*

gene, or gene product(s) (*e.g.* mRNA, lipin protein, *etc.*) provide good targets for new agents that modulate adipocyte differentiation and/or modulate fat accumulation. Thus, in one embodiment, this invention provides methods of screening for agents that modulate *Lpin1* expression and/or activity and hence adipocyte differentiation and/or fat

5 accumulation. The methods preferably involve detecting a change in the expression level and/or activity level of a *Lpin1* gene or gene product (*e.g.* Lpin protein) in cell(s) contacted with the test agent in question. An elevated *Lpin1* expression level or activity level in the presence of the agent, *e.g.*, as compared to a negative control where the test agent is absent or at reduced concentration indicates that the agent upregulates *Lpin1* activity or
10 expression. Conversely, decreased *Lpin1* expression level or activity level in the presence of the agent as compared to a negative control where the test agent is absent or at reduced concentration indicates that the agent down-regulates *Lpin1* activity or expression.

[0042] Expression levels of a gene can be altered by changes in the transcription of the gene product (*i.e.* transcription of mRNA), and/or by changes in translation of the gene
15 product (*i.e.* translation of the protein), and/or by post-translational modification(s) (*e.g.* protein folding, glycosylation, *etc.*). Thus preferred assays of this invention include assaying for level of transcribed mRNA (or other nucleic acids derived from the *Lpin1* gene), level of translated protein, activity of translated protein, *etc.* Examples of such approaches are described below.

20 A) Nucleic-acid based assays.

1) Target molecules.

[0043] Changes in expression level can be detected by measuring changes in mRNA and/or a nucleic acid derived from the mRNA (*e.g.* reverse-transcribed cDNA, *etc.*). In order to measure the *Lpin1* expression level it is desirable to provide a nucleic
25 acid sample for such analysis. In preferred embodiments the nucleic acid is found in or derived from a biological sample. The term "biological sample", as used herein, refers to a sample obtained from an organism or from components (*e.g.*, cells) of an organism. The sample may be of any biological tissue or fluid. Biological samples may also include organs or sections of tissues such as frozen sections taken for histological purposes.

[0044] The nucleic acid (*e.g.*, mRNA nucleic acid derived from mRNA) is, in certain preferred embodiments, isolated from the sample according to any of a number of methods well known to those of skill in the art. Methods of isolating mRNA are well known to those of skill in the art. For example, methods of isolation and purification of nucleic acids are described in detail in by Tijssen ed., (1993) Chapter 3 of *Laboratory Techniques in Biochemistry and Molecular Biology: Hybridization With Nucleic Acid Probes, Part I. Theory and Nucleic Acid Preparation*, Elsevier, N.Y. and Tijssen ed.

[0045] In a preferred embodiment, the "total" nucleic acid is isolated from a given sample using, for example, an acid guanidinium-phenol-chloroform extraction method and polyA⁺ mRNA is isolated by oligo dT column chromatography or by using (dT)_n magnetic beads (*see, e.g.*, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual* (2nd ed.), Vols. 1-3, Cold Spring Harbor Laboratory, (1989), or *Current Protocols in Molecular Biology*, F. Ausubel *et al.*, ed. Greene Publishing and Wiley-Interscience, New York (1987)).

[0046] Frequently, it is desirable to amplify the nucleic acid sample prior to assaying for expression level. Methods of amplifying nucleic acids are well known to those of skill in the art and include, but are not limited to polymerase chain reaction (PCR, *see, e.g.*, Innis, *et al.*, (1990) *PCR Protocols. A guide to Methods and Application*. Academic Press, Inc. San Diego,)), ligase chain reaction (LCR) (*see* Wu and Wallace (1989) *Genomics* 4: 560, Landegren *et al.* (1988) *Science* 241: 1077, and Barringer *et al.* (1990) *Gene* 89: 117, transcription amplification (Kwoh *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86: 1173), self-sustained sequence replication (Guatelli *et al.* (1990) *Proc. Nat. Acad. Sci. USA* 87: 1874), dot PCR, and linker adapter PCR, *etc.*).

[0047] In a particularly preferred embodiment, where it is desired to quantify the transcription level (and thereby expression) of *Lpin1* in a sample, the nucleic acid sample is one in which the concentration of the *Lpin1* mRNA transcript(s), or the concentration of the nucleic acids derived from the *Lpin1* mRNA transcript(s), is proportional to the transcription level (and therefore expression level) of that gene. Similarly, it is preferred that the hybridization signal intensity be proportional to the amount of hybridized nucleic acid. While it is preferred that the proportionality be relatively strict (*e.g.*, a doubling in transcription rate results in a doubling in mRNA transcript in the sample nucleic acid pool

and a doubling in hybridization signal), one of skill will appreciate that the proportionality can be more relaxed and even non-linear. Thus, for example, an assay where a 5 fold difference in concentration of the target mRNA results in a 3 to 6 fold difference in hybridization intensity is sufficient for most purposes.

5 [0048] Where more precise quantification is required appropriate controls can be run to correct for variations introduced in sample preparation and hybridization as described herein. In addition, serial dilutions of "standard" target nucleic acids (*e.g.*, mRNAs) can be used to prepare calibration curves according to methods well known to those of skill in the art. Of course, where simple detection of the presence or absence of a
10 transcript or large differences of changes in nucleic acid concentration is desired, no elaborate control or calibration is required.

[0049] In the simplest embodiment, the *Lpin1*-containing nucleic acid sample is the total mRNA or a total cDNA isolated and/or otherwise derived from a biological sample. The nucleic acid may be isolated from the sample according to any of a number
15 of methods well known to those of skill in the art as indicated above.

2) Hybridization-based assays.

[0050] Using the *Lpin1* sequences provided herein (*see, e.g.*, SEQ ID Nos: 1 and 2) detecting and/or quantifying the *Lpin1* transcript(s) can be routinely accomplished using nucleic acid hybridization techniques (*see, e.g.*, Sambrook *et al. supra*). For example, one
20 method for evaluating the presence, absence, or quantity of *Lpin1* reverse-transcribed cDNA involves a "Southern Blot". In a Southern Blot, the DNA (*e.g.*, reverse-transcribed *Lpin1* mRNA), typically fragmented and separated on an electrophoretic gel, is hybridized to a probe specific for *Lpin1* (or to a mutant thereof). Comparison of the intensity of the hybridization signal from the *Lpin1* probe with a "control" probe (*e.g.* a probe for a
25 "housekeeping gene) provides an estimate of the relative expression level of the target nucleic acid.

[0051] Alternatively, the *Lpin1* mRNA can be directly quantified in a Northern blot. In brief, the mRNA is isolated from a given cell sample using, for example, an acid guanidinium-phenol-chloroform extraction method. The mRNA is then electrophoresed to
30 separate the mRNA species and the mRNA is transferred from the gel to a nitrocellulose

membrane. As with the Southern blots, labeled probes are used to identify and/or quantify the target *Lpin1* mRNA. Appropriate controls (e.g. probes to housekeeping genes) provide a reference for evaluating relative expression level.

[0052] An alternative means for determining the *Lpin1* expression level is *in situ* hybridization. *In situ* hybridization assays are well known (e.g., Angerer (1987) *Meth. Enzymol* 152: 649). Generally, *in situ* hybridization comprises the following major steps: (1) fixation of tissue or biological structure to be analyzed; (2) prehybridization treatment of the biological structure to increase accessibility of target DNA, and to reduce nonspecific binding; (3) hybridization of the mixture of nucleic acids to the nucleic acid in the biological structure or tissue; (4) post-hybridization washes to remove nucleic acid fragments not bound in the hybridization and (5) detection of the hybridized nucleic acid fragments. The reagent used in each of these steps and the conditions for use vary depending on the particular application.

[0053] In some applications it is necessary to block the hybridization capacity of repetitive sequences. Thus, in some embodiments, tRNA, human genomic DNA, or Cot-1 DNA is used to block non-specific hybridization.

3) Amplification-based assays.

[0054] In another embodiment, amplification-based assays can be used to measure *Lpin1* expression (transcription) level. In such amplification-based assays, the target nucleic acid sequences (i.e., *Lpin1*) act as template(s) in amplification reaction(s) (e.g. Polymerase Chain Reaction (PCR) or reverse-transcription PCR (RT-PCR)). In a quantitative amplification, the amount of amplification product will be proportional to the amount of template (e.g., *Lpin1* mRNA) in the original sample. Comparison to appropriate (e.g. healthy tissue or cells unexposed to the test agent) controls provides a measure of the *Lpin1* transcript level.

[0055] Methods of "quantitative" amplification are well known to those of skill in the art. For example, quantitative PCR involves simultaneously co-amplifying a known quantity of a control sequence using the same primers. This provides an internal standard that may be used to calibrate the PCR reaction. Detailed protocols for quantitative PCR are provided in Innis *et al.* (1990) *PCR Protocols, A Guide to Methods and Applications*,

Academic Press, Inc. N.Y.). One approach, for example, involves simultaneously co-amplifying a known quantity of a control sequence using the same primers as those used to amplify the target. This provides an internal standard that may be used to calibrate the PCR reaction.

- 5 [0056] One preferred internal standard is a synthetic AW106 cRNA. The AW106 cRNA is combined with RNA isolated from the sample according to standard techniques known to those of skill in the art. The RNA is then reverse transcribed using a reverse transcriptase to provide copy DNA. The cDNA sequences are then amplified (*e.g.*, by
- 10 electrophoresis, and the amount of labeled nucleic acid (proportional to the amount of amplified product) is determined. The amount of mRNA in the sample is then calculated by comparison with the signal produced by the known AW106 RNA standard. Detailed protocols for quantitative PCR are provided in PCR Protocols, A Guide to Methods and Applications, Innis *et al.* (1990) Academic Press, Inc. N.Y.. The known nucleic acid
- 15 sequence(s) for *Lpin1* are sufficient to enable one of skill to routinely select primers to amplify any portion of the gene.

4) Hybridization Formats and Optimization of hybridization conditions.

a) Array-based hybridization formats.

- 20 [0057] In one embodiment, the methods of this invention can be utilized in array-based hybridization formats. Arrays are a multiplicity of different "probe" or "target" nucleic acids (or other compounds) attached to one or more surfaces (*e.g.*, solid, membrane, or gel). In a preferred embodiment, the multiplicity of nucleic acids (or other moieties) is attached to a single contiguous surface or to a multiplicity of surfaces
- 25 juxtaposed to each other.

- [0058] In an array format a large number of different hybridization reactions can be run essentially "in parallel." This provides rapid, essentially simultaneous, evaluation of a number of hybridizations in a single "experiment". Methods of performing hybridization reactions in array based formats are well known to those of skill in the art
- 30 (*see, e.g.*, Pastinen (1997) *Genome Res.* 7: 606-614; Jackson (1996) *Nature Biotechnology*

14:1685; Chee (1995) *Science* 274: 610; WO 96/17958, Pinkel *et al.* (1998) *Nature Genetics* 20: 207-211).

[0059] Arrays, particularly nucleic acid arrays can be produced according to a wide variety of methods well known to those of skill in the art. For example, in a simple embodiment, "low density" arrays can simply be produced by spotting (*e.g.* by hand using a pipette) different nucleic acids at different locations on a solid support (*e.g.* a glass surface, a membrane, *etc.*).

[0060] This simple spotting, approach has been automated to produce high density spotted arrays (*see, e.g.*, U.S. Patent No: 5,807,522). This patent describes the use of an automated system that taps a microcapillary against a surface to deposit a small volume of a biological sample. The process is repeated to generate high-density arrays.

[0061] Arrays can also be produced using oligonucleotide synthesis technology. Thus, for example, U.S. Patent No. 5,143,854 and PCT Patent Publication Nos. WO 90/15070 and 92/10092 teach the use of light-directed combinatorial synthesis of high density oligonucleotide arrays. Synthesis of high-density arrays is also described in U.S. Patents 5,744,305, 5,800,992 and 5,445,934.

b) Other hybridization formats.

[0062] As indicated above a variety of nucleic acid hybridization formats are known to those skilled in the art. For example, common formats include sandwich assays and competition or displacement assays. Such assay formats are generally described in Hames and Higgins (1985) *Nucleic Acid Hybridization, A Practical Approach*, IRL Press; Gall and Pardue (1969) *Proc. Natl. Acad. Sci. USA* 63: 378-383; and John *et al.* (1969) *Nature* 223: 582-587.

[0063] Sandwich assays are commercially useful hybridization assays for detecting or isolating nucleic acid sequences. Such assays utilize a "capture" nucleic acid covalently immobilized to a solid support and a labeled "signal" nucleic acid in solution. The sample will provide the target nucleic acid. The "capture" nucleic acid and "signal" nucleic acid probe hybridize with the target nucleic acid to form a "sandwich" hybridization complex. To be most effective, the signal nucleic acid should not hybridize with the capture nucleic acid.

[0064] Typically, labeled signal nucleic acids are used to detect hybridization. Complementary nucleic acids or signal nucleic acids may be labeled by any one of several methods typically used to detect the presence of hybridized polynucleotides. The most common method of detection is the use of autoradiography with ^3H , ^{125}I , ^{35}S , ^{14}C , or ^{32}P -labelled probes or the like. Other labels include ligands that bind to labeled antibodies, fluorophores, chemiluminescent agents, enzymes, and antibodies that can serve as specific binding pair members for a labeled ligand.

[0065] Detection of a hybridization complex may require the binding of a signal generating complex to a duplex of target and probe polynucleotides or nucleic acids.

Typically, such binding occurs through ligand and anti-ligand interactions as between a ligand-conjugated probe and an anti-ligand conjugated with a signal.

[0066] The sensitivity of the hybridization assays may be enhanced through use of a nucleic acid amplification system that multiplies the target nucleic acid being detected. Examples of such systems include the polymerase chain reaction (PCR) system and the ligase chain reaction (LCR) system. Other methods recently described in the art are the nucleic acid sequence based amplification (NASBAO, Cangene, Mississauga, Ontario) and Q Beta Replicase systems.

c) Optimization of hybridization conditions.

[0067] Nucleic acid hybridization simply involves providing a denatured probe and target nucleic acid under conditions where the probe and its complementary target can form stable hybrid duplexes through complementary base pairing. The nucleic acids that do not form hybrid duplexes are then washed away leaving the hybridized nucleic acids to be detected, typically through detection of an attached detectable label. It is generally recognized that nucleic acids are denatured by increasing the temperature or decreasing the salt concentration of the buffer containing the nucleic acids, or in the addition of chemical agents, or the raising of the pH. Under low stringency conditions (*e.g.*, low temperature and/or high salt and/or high target concentration) hybrid duplexes (*e.g.*, DNA:DNA, RNA:RNA, or RNA:DNA) will form even where the annealed sequences are not perfectly complementary. Thus specificity of hybridization is reduced at lower stringency. Conversely, at higher stringency (*e.g.*, higher temperature or lower salt) successful hybridization requires fewer mismatches.

[0068] One of skill in the art will appreciate that hybridization conditions may be selected to provide any degree of stringency. In a preferred embodiment, hybridization is performed at low stringency to ensure hybridization and then subsequent washes are performed at higher stringency to eliminate mismatched hybrid duplexes. Successive washes may be performed at increasingly higher stringency (*e.g.*, down to as low as 0.25 X SSPE at 37°C to 70°C) until a desired level of hybridization specificity is obtained. Stringency can also be increased by addition of agents such as formamide. Hybridization specificity may be evaluated by comparison of hybridization to the test probes with hybridization to the various controls that can be present.

[0069] In general, there is a tradeoff between hybridization specificity (stringency) and signal intensity. Thus, in a preferred embodiment, the wash is performed at the highest stringency that produces consistent results and that provides a signal intensity greater than approximately 10% of the background intensity. Thus, in a preferred embodiment, the hybridized array may be washed at successively higher stringency solutions and read between each wash. Analysis of the data sets thus produced will reveal a wash stringency above which the hybridization pattern is not appreciably altered and which provides adequate signal for the particular probes of interest.

[0070] In a preferred embodiment, background signal is reduced by the use of a blocking reagent (*e.g.*, tRNA, sperm DNA, cot-1 DNA, *etc.*) during the hybridization to reduce non-specific binding. The use of blocking agents in hybridization is well known to those of skill in the art (*see, e.g.*, Chapter 8 in P. Tijssen, *supra.*)

[0071] Methods of optimizing hybridization conditions are well known to those of skill in the art (*see, e.g.*, Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology, Vol. 24: Hybridization With Nucleic Acid Probes*, Elsevier, N.Y.).

[0072] Optimal conditions are also a function of the sensitivity of label (*e.g.*, fluorescence) detection for different combinations of substrate type, fluorochrome, excitation and emission bands, spot size and the like. Low fluorescence background surfaces can be used (*see, e.g.*, Chu (1992) *Electrophoresis* 13:105-114). The sensitivity for detection of spots ("target elements") of various diameters on the candidate surfaces can be readily determined by, *e.g.*, spotting a dilution series of fluorescently end labeled DNA fragments. These spots are then imaged using conventional fluorescence

microscopy. The sensitivity, linearity, and dynamic range achievable from the various combinations of fluorochrome and solid surfaces (*e.g.*, glass, fused silica, *etc.*) can thus be determined. Serial dilutions of pairs of fluorochrome in known relative proportions can also be analyzed. This determines the accuracy with which fluorescence ratio

5 measurements reflect actual fluorochrome ratios over the dynamic range permitted by the detectors and fluorescence of the substrate upon which the probe has been fixed.

d) Labeling and detection of nucleic acids.

[0073] The probes used herein for detection of *Lpin1* expression levels can be full length or less than the full length of the *Lpin1* or mutants thereof. Shorter probes are

10 empirically tested for specificity. Preferred probes are sufficiently long so as to specifically hybridize with the *Lpin1* target nucleic acid(s) under stringent conditions. The preferred size range is from about 20 bases to the length of the *Lpin1* mRNA, more preferably from about 30 bases to the length of the *Lpin1* mRNA, and most preferably from about 40 bases to the length of the *Lpin1* mRNA.

15 [0074] The probes are typically labeled, with a detectable label. Detectable labels suitable for use in the present invention include any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, electrical, optical or chemical means. Useful labels in the present invention include biotin for staining with labeled streptavidin conjugate, magnetic beads (*e.g.*, DynabeadsTM), fluorescent dyes (*e.g.*,

20 fluorescein, texas red, rhodamine, green fluorescent protein, and the like, *see, e.g.*, Molecular Probes, Eugene, Oregon, USA), radiolabels (*e.g.*, ³H, ¹²⁵I, ³⁵S, ¹⁴C, or ³²P), enzymes (*e.g.*, horse radish peroxidase, alkaline phosphatase and others commonly used in an ELISA), and colorimetric labels such as colloidal gold (*e.g.*, gold particles in the 40 -80

25 (*e.g.*, polystyrene, polypropylene, latex, *etc.*) beads. Patents teaching the use of such labels include U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241.

[0075] A fluorescent label is preferred because it provides a very strong signal with low background. It is also optically detectable at high resolution and sensitivity

30 through a quick scanning procedure. The nucleic acid samples can all be labeled with a single label, *e.g.*, a single fluorescent label. Alternatively, in another embodiment,

different nucleic acid samples can be simultaneously hybridized where each nucleic acid sample has a different label. For instance, one target could have a green fluorescent label and a second target could have a red fluorescent label. The scanning step will distinguish sites of binding of the red label from those binding the green fluorescent label. Each

5 nucleic acid sample (target nucleic acid) can be analyzed independently from one another.

[0076] Suitable chromogens which can be employed include those molecules and compounds which absorb light in a distinctive range of wavelengths so that a color can be observed or, alternatively, which emit light when irradiated with radiation of a particular wave length or wave length range, *e.g.*, fluorescers.

10 [0077] Desirably, fluorescent labels should absorb light above about 300 nm, preferably about 350 nm, and more preferably above about 400 nm, usually emitting at wavelengths greater than about 10 nm higher than the wavelength of the light absorbed. It should be noted that the absorption and emission characteristics of the bound dye can differ from the unbound dye. Therefore, when referring to the various wavelength ranges

15 and characteristics of the dyes, it is intended to indicate the dyes as employed and not the dye which is unconjugated and characterized in an arbitrary solvent.

[0078] Detectable signal can also be provided by chemiluminescent and bioluminescent sources. Chemiluminescent sources include a compound which becomes electronically excited by a chemical reaction and can then emit light which serves as the

20 detectable signal or donates energy to a fluorescent acceptor. Alternatively, luciferins can be used in conjunction with luciferase or lucigenins to provide bioluminescence.

[0079] Spin labels are provided by reporter molecules with an unpaired electron spin which can be detected by electron spin resonance (ESR) spectroscopy. Exemplary spin labels include organic free radicals, transitional metal complexes, particularly

25 vanadium, copper, iron, and manganese, and the like. Exemplary spin labels include nitroxide free radicals.

[0080] The label can be added to the target (sample) nucleic acid(s) prior to, or after the hybridization. So called "direct labels" are detectable labels that are directly attached to or incorporated into the target (sample) nucleic acid prior to hybridization. In

30 contrast, so called "indirect labels" are joined to the hybrid duplex after hybridization. Often, the indirect label is attached to a binding moiety that has been attached to the target

nucleic acid prior to the hybridization. Thus, for example, the target nucleic acid may be biotinylated before the hybridization. After hybridization, an avidin-conjugated fluorophore will bind the biotin bearing hybrid duplexes providing a label that is easily detected. For a detailed review of methods of labeling nucleic acids and detecting labeled hybridized nucleic acids see *Laboratory Techniques in Biochemistry and Molecular Biology, Vol. 24: Hybridization With Nucleic Acid Probes*, P. Tijssen, ed. Elsevier, N.Y., (1993)).

[0081] Fluorescent labels are easily added during an *in vitro* transcription reaction. Thus, for example, fluorescein labeled UTP and CTP can be incorporated into the RNA produced in an *in vitro* transcription.

[0082] The labels can be attached directly or through a linker moiety. In general, the site of label or linker-label attachment is not limited to any specific position. For example, a label may be attached to a nucleoside, nucleotide, or analogue thereof at any position that does not interfere with detection or hybridization as desired. For example, certain Label-ON Reagents from Clontech (Palo Alto, CA) provide for labeling interspersed throughout the phosphate backbone of an oligonucleotide and for terminal labeling at the 3' and 5' ends. As shown for example herein, labels can be attached at positions on the ribose ring or the ribose can be modified and even eliminated as desired. The base moieties of useful labeling reagents can include those that are naturally occurring or modified in a manner that does not interfere with the purpose to which they are put. Modified bases include but are not limited to 7-deaza A and G, 7-deaza-8-aza A and G, and other heterocyclic moieties.

[0083] It will be recognized that fluorescent labels are not to be limited to single species organic molecules, but include inorganic molecules, multi-molecular mixtures of organic and/or inorganic molecules, crystals, heteropolymers, and the like. Thus, for example, CdSe-CdS core-shell nanocrystals enclosed in a silica shell can be easily derivatized for coupling to a biological molecule (Bruchez *et al.* (1998) *Science*, 281: 2013-2016). Similarly, highly fluorescent quantum dots (zinc sulfide-capped cadmium selenide) have been covalently coupled to biomolecules for use in ultrasensitive biological detection (Warren and Nie (1998) *Science*, 281: 2016-2018).

B) Polypeptide-based assays.**1) Assay Formats.**

[0084] In addition to, or in alternative to, the detection of *Lpin1* nucleic acid expression level(s), alterations in expression of lipin can be detected and/or quantified by detecting and/or quantifying the amount and/or activity of translated lipin polypeptide.

2) Detection of expressed protein

[0085] The polypeptide(s) encoded by the *Lpin1* gene can be detected and quantified by any of a number of methods well known to those of skill in the art. These may include analytic biochemical methods such as electrophoresis, capillary electrophoresis, high performance liquid chromatography (HPLC), thin layer chromatography (TLC), hyperdiffusion chromatography, and the like, or various immunological methods such as fluid or gel precipitin reactions, immunodiffusion (single or double), immunoelectrophoresis, radioimmunoassay (RIA), enzyme-linked immunosorbent assays (ELISAs), immunofluorescent assays, western blotting, and the like.

[0086] In one preferred embodiment, the lipin polypeptide(s) are detected/quantified in an electrophoretic protein separation (e.g. a 1- or 2-dimensional electrophoresis). Means of detecting proteins using electrophoretic techniques are well known to those of skill in the art (*see generally*, R. Scopes (1982) *Protein Purification*, Springer-Verlag, N.Y.; Deutscher, (1990) *Methods in Enzymology Vol. 182: Guide to Protein Purification*, Academic Press, Inc., N.Y.).

[0087] In another preferred embodiment, Western blot (immunoblot) analysis is used to detect and quantify the presence of polypeptide(s) of this invention in the sample. This technique generally comprises separating sample proteins by gel electrophoresis on the basis of molecular weight, transferring the separated proteins to a suitable solid support, (such as a nitrocellulose filter, a nylon filter, or derivatized nylon filter), and incubating the sample with the antibodies that specifically bind the target polypeptide(s).

[0088] The antibodies specifically bind to the target polypeptide(s) and may be directly labeled or alternatively may be subsequently detected using labeled antibodies

(e.g., labeled sheep anti-mouse antibodies) that specifically bind to a domain of the antibody.

[0089] In preferred embodiments, the lipin polypeptide(s) are detected using an immunoassay. As used herein, an immunoassay is an assay that utilizes an antibody to specifically bind to the analyte (e.g., the target polypeptide(s)). The immunoassay is thus characterized by detection of specific binding of a polypeptide of this invention to an antibody as opposed to the use of other physical or chemical properties to isolate, target, and quantify the analyte.

[0090] Any of a number of well recognized immunological binding assays (*see, e.g.,* U.S. Patents 4,366,241; 4,376,110; 4,517,288; and 4,837,168) are well suited to detection or quantification of the polypeptide(s) identified herein.. For a review of the general immunoassays, see also Asai (1993) *Methods in Cell Biology Volume 37: Antibodies in Cell Biology*, Academic Press, Inc. New York; Stites & Terr (1991) *Basic and Clinical Immunology 7th Edition*.

[0091] Immunological binding assays (or immunoassays) typically utilize a “capture agent” to specifically bind to and often immobilize the analyte (lipin polypeptide). In preferred embodiments, the capture agent is an antibody.

[0092] Immunoassays also often utilize a labeling agent to specifically bind to and label the binding complex formed by the capture agent and the analyte. The labeling agent may itself be one of the moieties comprising the antibody/analyte complex. Thus, the labeling agent may be a labeled polypeptide or a labeled antibody that specifically recognizes the already bound target polypeptide. Alternatively, the labeling agent may be a third moiety, such as another antibody, that specifically binds to the capture agent /polypeptide complex.

[0093] Other proteins capable of specifically binding immunoglobulin constant regions, such as protein A or protein G may also be used as the label agent. These proteins are normal constituents of the cell walls of streptococcal bacteria. They exhibit a strong non-immunogenic reactivity with immunoglobulin constant regions from a variety of species (*see, generally* Kronval, *et al.* (1973) *J. Immunol.*, 111: 1401-1406, and Akerstrom (1985) *J. Immunol.*, 135: 2589-2542).

[0094] Preferred immunoassays for detecting the target polypeptide(s) are either competitive or noncompetitive. Noncompetitive immunoassays are assays in which the amount of captured analyte is directly measured. In one preferred "sandwich" assay, for example, the capture agents (antibodies) can be bound directly to a solid substrate where they are immobilized. These immobilized antibodies then capture the target polypeptide present in the test sample. The target polypeptide thus immobilized is then bound by a labeling agent, such as a second antibody bearing a label.

[0095] In competitive assays, the amount of analyte (lipin polypeptide) present in the sample is measured indirectly by measuring the amount of an added (exogenous) analyte displaced (or competed away) from a capture agent (antibody) by the analyte present in the sample. In one competitive assay, a known amount of, in this case, labeled polypeptide is added to the sample and the sample is then contacted with a capture agent. The amount of labeled polypeptide bound to the antibody is inversely proportional to the concentration of target polypeptide present in the sample.

[0096] In one particularly preferred embodiment, the antibody is immobilized on a solid substrate. The amount of target polypeptide bound to the antibody may be determined either by measuring the amount of target polypeptide present in an polypeptide /antibody complex, or alternatively by measuring the amount of remaining uncomplexed polypeptide.

[0097] The immunoassay methods of the present invention include an enzyme immunoassay (EIA) which utilizes, depending on the particular protocol employed, unlabeled or labeled (*e.g.*, enzyme-labeled) derivatives of polyclonal or monoclonal antibodies or antibody fragments or single-chain antibodies that bind lipin polypeptide(s), either alone or in combination. In the case where the antibody that binds lipin polypeptide is not labeled, a different detectable marker, for example, an enzyme-labeled antibody capable of binding to the monoclonal antibody which binds the lipin polypeptide, may be employed. Any of the known modifications of EIA, for example, enzyme-linked immunoabsorbent assay (ELISA), may also be employed. As indicated above, also contemplated by the present invention are immunoblotting immunoassay techniques such as western blotting employing an enzymatic detection system.

[0098] The immunoassay methods of the present invention may also be other known immunoassay methods, for example, fluorescent immunoassays using antibody conjugates or antigen conjugates of fluorescent substances such as fluorescein or rhodamine, latex agglutination with antibody-coated or antigen-coated latex particles, 5 haemagglutination with antibody-coated or antigen-coated red blood corpuscles, and immunoassays employing an avidin-biotin or streptavidin-biotin detection systems, and the like.

[0099] The particular parameters employed in the immunoassays of the present invention can vary widely depending on various factors such as the concentration of antigen in the sample, the nature of the sample, the type of immunoassay employed and the like. Optimal conditions can be readily established by those of ordinary skill in the art. In certain embodiments, the amount of antibody that binds lipin polypeptides is typically selected to give 50% binding of detectable marker in the absence of sample. If purified antibody is used as the antibody source, the amount of antibody used per assay will 15 generally range from about 1 ng to about 100 ng. Typical assay conditions include a temperature range of about 4°C to about 45°C, preferably about 25°C to about 37°C, and most preferably about 25°C, a pH value range of about 5 to 9, preferably about 7, and an ionic strength varying from that of distilled water to that of about 0.2M sodium chloride, preferably about that of 0.15M sodium chloride. Times will vary widely depending upon the nature of the assay, and generally range from about 0.1 minute to about 24 hours. A 20 wide variety of buffers, for example PBS, may be employed, and other reagents such as salt to enhance ionic strength, proteins such as serum albumins, stabilizers, biocides and non-ionic detergents may also be included.

[0100] The assays of this invention are scored (as positive or negative or quantity of target polypeptide) according to standard methods well known to those of skill in the art. The particular method of scoring will depend on the assay format and choice of label. For example, a Western Blot assay can be scored by visualizing the colored product produced by the enzymatic label. A clearly visible colored band or spot at the correct molecular weight is scored as a positive result, while the absence of a clearly visible spot 25 or band is scored as a negative. The intensity of the band or spot can provide a 30 quantitative measure of target polypeptide concentration.

[0101] Antibodies for use in the various immunoassays described herein are commercially available or can be produced as described below.

3) Antibodies to lipin polypeptides.

[0102] Either polyclonal or monoclonal antibodies may be used in the immunoassays of the invention described herein. Polyclonal antibodies are preferably raised by multiple injections (*e.g.* subcutaneous or intramuscular injections) of substantially pure polypeptides or antigenic polypeptides into a suitable non-human mammal. The antigenicity of the target peptides can be determined by conventional techniques to determine the magnitude of the antibody response of an animal that has been immunized with the peptide. Generally, the peptides that are used to raise antibodies for use in the methods of this invention should generally be those which induce production of high titers of antibody with relatively high affinity for target polypeptides encoded by *Lpin1* or variants thereof.

[0103] If desired, the immunizing peptide may be coupled to a carrier protein by conjugation using techniques that are well-known in the art. Such commonly used carriers which are chemically coupled to the peptide include keyhole limpet hemocyanin (KLH), thyroglobulin, bovine serum albumin (BSA), and tetanus toxoid. The coupled peptide is then used to immunize the animal (*e.g.* a mouse or a rabbit).

[0104] The antibodies are then obtained from blood samples taken from the mammal. The techniques used to develop polyclonal antibodies are known in the art (see, *e.g.*, *Methods of Enzymology*, "Production of Antisera With Small Doses of Immunogen: Multiple Intradermal Injections", Langone, *et al.* eds. (Acad. Press, 1981)). Polyclonal antibodies produced by the animals can be further purified, for example, by binding to and elution from a matrix to which the peptide to which the antibodies were raised is bound. Those of skill in the art will know of various techniques common in the immunology arts for purification and/or concentration of polyclonal antibodies, as well as monoclonal antibodies *see*, for example, Coligan, *et al.* (1991) Unit 9, *Current Protocols in Immunology*, Wiley Interscience).

[0105] Preferably, however, the antibodies produced will be monoclonal antibodies ("mAb's"). For preparation of monoclonal antibodies, immunization of a

mouse or rat is preferred. The term "antibody" as used in this invention includes intact molecules as well as fragments thereof, such as, Fab and F(ab')², and/or single-chain antibodies (*e.g.* scFv) which are capable of binding an epitopic determinant. Also, in this context, the term "mab's of the invention" refers to monoclonal antibodies with specificity for a polypeptide encoded by *Lpin1*.

[0106] The general method used for production of hybridomas secreting mAbs is well known (Kohler and Milstein (1975) *Nature*, 256:495). Briefly, as described by Kohler and Milstein the technique comprised isolating lymphocytes from regional draining lymph nodes of five separate cancer patients with either melanoma, teratocarcinoma or cancer of the cervix, glioma or lung, (where samples were obtained from surgical specimens), pooling the cells, and fusing the cells with SHFP-1. Hybridomas were screened for production of antibody which bound to cancer cell lines. Confirmation of specificity among mAb's can be accomplished using relatively routine screening techniques (such as the enzyme-linked immunosorbent assay, or "ELISA") to determine the elementary reaction pattern of the mAb of interest.

[0107] Antibodies fragments, *e.g.* single chain antibodies (scFv or others), can also be produced/selected using phage display technology. The ability to express antibody fragments on the surface of viruses that infect bacteria (bacteriophage or phage) makes it possible to isolate a single binding antibody fragment, *e.g.*, from a library of greater than 10¹⁰ nonbinding clones. To express antibody fragments on the surface of phage (phage display), an antibody fragment gene is inserted into the gene encoding a phage surface protein (*e.g.*, pIII) and the antibody fragment-pIII fusion protein is displayed on the phage surface (McCafferty *et al.* (1990) *Nature*, 348: 552-554; Hoogenboom *et al.* (1991) *Nucleic Acids Res.* 19: 4133-4137).

[0108] Since the antibody fragments on the surface of the phage are functional, phage bearing antigen binding antibody fragments can be separated from non-binding phage by antigen affinity chromatography (McCafferty *et al.* (1990) *Nature*, 348: 552-554). Depending on the affinity of the antibody fragment, enrichment factors of 20 fold - 1,000,000 fold are obtained for a single round of affinity selection. By infecting bacteria with the eluted phage, however, more phage can be grown and subjected to another round of selection. In this way, an enrichment of 1000 fold in one round can become 1,000,000

fold in two rounds of selection (McCafferty *et al.* (1990) *Nature*, 348: 552-554). Thus even when enrichments are low (Marks *et al.* (1991) *J. Mol. Biol.* 222: 581-597), multiple rounds of affinity selection can lead to the isolation of rare phage. Since selection of the phage antibody library on antigen results in enrichment, the majority of clones bind antigen after as few as three to four rounds of selection. Thus only a relatively small number of clones (several hundred) need to be analyzed for binding to antigen.

[0109] Human antibodies can be produced without prior immunization by displaying very large and diverse V-gene repertoires on phage (Marks *et al.* (1991) *J. Mol. Biol.* 222: 581-597). In one embodiment natural V_H and V_L repertoires present in human peripheral blood lymphocytes are were isolated from unimmunized donors by PCR. The V-gene repertoires were spliced together at random using PCR to create a scFv gene repertoire which is was cloned into a phage vector to create a library of 30 million phage antibodies (*Id.*). From this single "naive" phage antibody library, binding antibody fragments have been isolated against more than 17 different antigens, including haptens, polysaccharides and proteins (Marks *et al.* (1991) *J. Mol. Biol.* 222: 581-597; Marks *et al.* (1993). *Bio/Technology*. 10: 779-783; Griffiths *et al.* (1993) *EMBO J.* 12: 725-734; Clackson *et al.* (1991) *Nature*. 352: 624-628). Antibodies have been produced against self proteins, including human thyroglobulin, immunoglobulin, tumor necrosis factor and CEA (Griffiths *et al.* (1993) *EMBO J.* 12: 725-734). It is also possible to isolate antibodies against cell surface antigens by selecting directly on intact cells. The antibody fragments are highly specific for the antigen used for selection and have affinities in the 1 :M to 100 nM range (Marks *et al.* (1991) *J. Mol. Biol.* 222: 581-597; Griffiths *et al.* (1993) *EMBO J.* 12: 725-734). Larger phage antibody libraries result in the isolation of more antibodies of higher binding affinity to a greater proportion of antigens.

[0110] It will also be recognized that antibodies can be prepared by any of a number of commercial services (*e.g.*, Berkeley antibody laboratories, Bethyl Laboratories, Anawa, Eurogenetec, *etc.*).

C) Assay Optimization.

[0111] The assays of this invention have immediate utility in screening for agents that modulate the *Lpin1* expression and/or activity in a cell, tissue or organism. The

assays of this invention can be optimized for use in particular contexts, depending, for example, on the source and/or nature of the biological sample and/or the particular test agents, and/or the analytic facilities available. Thus, for example, optimization can involve determining optimal conditions for binding assays, optimum sample processing conditions (e.g. preferred PCR conditions), hybridization conditions that maximize signal to noise, protocols that improve throughput, etc. In addition, assay formats can be selected and/or optimized according to the availability of equipment and/or reagents. Thus, for example, where commercial antibodies or ELISA kits are available it may be desired to assay protein concentration. Conversely, where it is desired to screen for modulators that alter transcription the *Lpin1* gene, nucleic acid based assays are preferred.

[0112] Routine selection and optimization of assay formats is well known to those of ordinary skill in the art.

D) Pre-screening for agents that bind *Lpin1* or lipin polypeptide

[0113] In certain embodiments it is desired to pre-screen test agents for the ability to interact with (e.g. specifically bind to) a *Lpin1* (or mutant/allele) nucleic acid or polypeptide. Specifically, binding test agents are more likely to interact with and thereby modulate *Lpin1* expression and/or activity. Thus, in some preferred embodiments, the test agent(s) are pre-screened for binding to *Lpin1* nucleic acids or to lipin proteins before performing the more complex assays described above.

[0114] In one embodiment, such pre-screening is accomplished with simple binding assays. Means of assaying for specific binding or the binding affinity of a particular ligand for a nucleic acid or for a protein are well known to those of skill in the art. In preferred binding assays, the lipin protein or nucleic acid is immobilized and exposed to a test agent (which can be labeled), or alternatively, the test agent(s) are immobilized and exposed to an lipin protein or to a *Lpin1* nucleic acid (which can be labeled). The immobilized moiety is then washed to remove any unbound material and the bound test agent or bound *Lpin1* nucleic acid or protein is detected (e.g. by detection of a label attached to the bound molecule). The amount of immobilized label is proportional to the degree of binding between the lipin protein or nucleic acid and the test agent.

E) Scoring the assay(s).

[0115] The assays of this invention are scored according to standard methods well known to those of skill in the art. The assays of this invention are typically scored as positive where there is a difference between the activity seen with the test agent present or where the test agent has been previously applied, and the (usually negative) control, preferably where the difference is statistically significant (*e.g.* at greater than 80%, preferably greater than about 90%, more preferably greater than about 98%, and most preferably greater than about 99% confidence level). Most preferred "positive" assays show at least a 1.2 fold, preferably at least a 1.5 fold, more preferably at least a 2 fold, and most preferably at least a 4 fold or even a 10-fold difference from the negative control.

F) Agents for screening: Combinatorial libraries (*e.g.*, small organic molecules).

[0116] Virtually any agent can be screened according to the methods of this invention. Such agents include, but are not limited to nucleic acids, proteins, sugars, polysaccharides, glycoproteins, lipids, and small organic molecules. The term small organic molecules typically refers to molecules of a size comparable to those organic molecules generally used in pharmaceuticals. The term excludes biological macromolecules (*e.g.*, proteins, nucleic acids, *etc.*). Preferred small organic molecules range in size up to about 5000 Da, more preferably up to 2000 Da, and most preferably up to about 1000 Da.

[0117] Conventionally, new chemical entities with useful properties are generated by identifying a chemical compound (called a "lead compound") with some desirable property or activity, creating variants of the lead compound, and evaluating the property and activity of those variant compounds. However, the current trend is to shorten the time scale for all aspects of drug discovery. Because of the ability to test large numbers quickly and efficiently, high throughput screening (HTS) methods are replacing conventional lead compound identification methods.

[0118] In one preferred embodiment, high throughput screening methods involve providing a library containing a large number of potential therapeutic compounds (candidate compounds). Such "combinatorial chemical libraries" are then screened in one

or more assays, as described herein to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as conventional "lead compounds" or can themselves be used as potential or actual therapeutics.

5 [0119] A combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis by combining a number of chemical "building blocks" such as reagents. For example, a linear combinatorial chemical library such as a polypeptide (*e.g.*, mutein) library is formed by combining a set of chemical building blocks called amino acids in every possible way for
10 a given compound length (*i.e.*, the number of amino acids in a polypeptide compound). Millions of chemical compounds can be synthesized through such combinatorial mixing of chemical building blocks. For example, one commentator has observed that the systematic, combinatorial mixing of 100 interchangeable chemical building blocks results in the theoretical synthesis of 100 million tetrameric compounds or 10 billion pentameric
15 compounds (Gallop *et al.* (1994) *J. Med. Chem.*, 37(9): 1233-1250).

[0120] Preparation of combinatorial chemical libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries (*see, e.g.*, U.S. Patent 5,010,175, Furka (1991) *Int. J. Pept. Prot. Res.*, 37: 487-493, Houghton *et al.* (1991) *Nature*, 354: 84-88). Peptide synthesis is by no means
20 the only approach envisioned and intended for use with the present invention. Other chemistries for generating chemical diversity libraries can also be used. Such chemistries include, but are not limited to: peptoids (PCT Publication No WO 91/19735, 26 Dec. 1991), encoded peptides (PCT Publication WO 93/20242, 14 Oct. 1993), random bio-oligomers (PCT Publication WO 92/00091, 9 Jan. 1992), benzodiazepines (U.S. Pat. No.
25 5,288,514), diversomers such as hydantoins, benzodiazepines and dipeptides (Hobbs *et al.*, (1993) *Proc. Nat. Acad. Sci. USA* 90: 6909-6913), vinylogous polypeptides (Hagihara *et al.* (1992) *J. Amer. Chem. Soc.* 114: 6568), nonpeptidal peptidomimetics with a Beta-D-Glucose scaffolding (Hirschmann *et al.*, (1992) *J. Amer. Chem. Soc.* 114: 9217-9218), analogous organic syntheses of small compound libraries (Chen *et al.* (1994) *J. Amer.*
30 *Chem. Soc.* 116: 2661), oligocarbamates (Cho, *et al.*, (1993) *Science* 261:1303), and/or peptidyl phosphonates (Campbell *et al.*, (1994) *J. Org. Chem.* 59: 658). *See, generally*, Gordon *et al.*, (1994) *J. Med. Chem.* 37:1385, nucleic acid libraries (*see, e.g.*, Strategene,

Corp.), peptide nucleic acid libraries (*see, e.g.*, U.S. Patent 5,539,083) antibody libraries (*see, e.g.*, Vaughn *et al.* (1996) *Nature Biotechnology*, 14(3): 309-314), and PCT/US96/10287), carbohydrate libraries (*see, e.g.*, Liang *et al.* (1996) *Science*, 274: 1520-1522, and U.S. Patent 5,593,853), and small organic molecule libraries (*see, e.g.*, benzodiazepines, Baum (1993) C&EN, Jan 18, page 33, isoprenoids U.S. Patent 5,569,588, thiazolidinones and metathiazanones U.S. Patent 5,549,974, pyrrolidines U.S. Patents 5,525,735 and 5,519,134, morpholino compounds U.S. Patent 5,506,337, benzodiazepines 5,288,514, and the like).

[0121] Devices for the preparation of combinatorial libraries are commercially available (*see, e.g.*, 357 MPS, 390 MPS, Advanced Chem Tech, Louisville KY, Symphony, Rainin, Woburn, MA, 433A Applied Biosystems, Foster City, CA, 9050 Plus, Millipore, Bedford, MA).

[0122] A number of well known robotic systems have also been developed for solution phase chemistries. These systems include, but are not limited to, automated workstations like the automated synthesis apparatus developed by Takeda Chemical Industries, LTD. (Osaka, Japan) and many robotic systems utilizing robotic arms (Zymate II, Zymark Corporation, Hopkinton, Mass.; Orca, Hewlett-Packard, Palo Alto, Calif.) which mimic the manual synthetic operations performed by a chemist and the VentureTM platform, an ultra-high-throughput synthesizer that can run between 576 and 9,600 simultaneous reactions from start to finish (*see* Advanced ChemTech, Inc. Louisville, KY)). Any of the above devices are suitable for use with the present invention. The nature and implementation of modifications to these devices (if any) so that they can operate as discussed herein will be apparent to persons skilled in the relevant art. In addition, numerous combinatorial libraries are themselves commercially available (*see, e.g.*, ComGenex, Princeton, N.J., Asinex, Moscow, Ru, Tripos, Inc., St. Louis, MO, ChemStar, Ltd, Moscow, RU, 3D Pharmaceuticals, Exton, PA, Martek Biosciences, Columbia, MD, *etc.*).

G) High Throughput Screening

[0123] Any of the assays for compounds modulating the accumulation or degradation of metabolic products described herein are amenable to high throughput

screening. Preferred assays detect increases or decreases in *Lpin1* transcription and/or translation in response to the presence of a test compound.

[0124] The cells utilized in the methods of this invention need not be contacted with a single test agent at a time. To the contrary, to facilitate high-throughput screening, a single cell may be contacted by at least two, preferably by at least 5, more preferably by at least 10, and most preferably by at least 20 test compounds. If the cell scores positive, it can be subsequently tested with a subset of the test agents until the agents having the activity are identified.

[0125] High throughput assays for various reporter gene products are well known to those of skill in the art. For example, multi-well fluorimeters are commercially available (*e.g.*, from Perkin-Elmer).

[0126] In addition, high throughput screening systems are commercially available (*see, e.g.*, Zymark Corp., Hopkinton, MA; Air Technical Industries, Mentor, OH; Beckman Instruments, Inc. Fullerton, CA; Precision Systems, Inc., Natick, MA, *etc.*).

These systems typically automate entire procedures including all sample and reagent pipetting, liquid dispensing, timed incubations, and final readings of the microplate in detector(s) appropriate for the assay. These configurable systems provide high throughput and rapid start up as well as a high degree of flexibility and customization. The manufacturers of such systems provide detailed protocols the various high throughput.

Thus, for example, Zymark Corp. provides technical bulletins describing screening systems for detecting the modulation of gene transcription, ligand binding, and the like.

H) Modulator databases.

[0127] In certain embodiments, the agents that score positively in the assays described herein (*e.g.* show an ability to modulate *Lpin1* expression) can be entered into a database of putative and/or actual modulators of *Lpin1* expression. The term database refers to a means for recording and retrieving information. In preferred embodiments the database also provides means for sorting and/or searching the stored information. The database can comprise any convenient media including, but not limited to, paper systems, card systems, mechanical systems, electronic systems, optical systems, magnetic systems or combinations thereof. Preferred databases include electronic (*e.g.* computer-based)

databases. Computer systems for use in storage and manipulation of databases are well known to those of skill in the art and include, but are not limited to "personal computer systems", mainframe systems, distributed nodes on an inter- or intra-net, data or databases stored in specialized hardware (*e.g.* in microchips), and the like.

5 **II. Identification of mutations in the LPIN1 gene in human lipodystrophy, obesity, diabetes, and atherosclerosis.**

[0128] In another embodiment, this invention provides methods of detecting a mutation (*e.g.* insertion, deletion, point mutation, *etc.*) in a *LPIN1* nucleic acid or in a polypeptide expressed from a *LPIN1* nucleic acid. Identification of such mutations
10 identifies/characterizes a lipodystrophic phenotype or a susceptibility to or likelihood of developing lipodystrophy, obesity, diabetes, atherosclerosis, and related pathologies. In other preferred embodiments, the methods involve detecting a *LPIN1* gene product (*e.g.* mRNA, polypeptide, *etc.*) where an alteration in the amount or activity of *LPIN1* gene product from the organism indicates that the organism has, or is susceptible to, a
15 lipodystrophic phenotype, obesity, diabetes or atherosclerosis.

[0129] Expression levels of a *LPIN1* gene product can be determined according to standard methods well known to those of skill in the art (*see, e.g.,* description of methods provided above). Similarly, mutations in *LPIN1* nucleic acids and/or lipin polypeptides can be identified according to standard methods. Thus, for example, mutations in nucleic
20 acids can be readily identified using single-probe and/or array-based hybridization methods (*see, e.g.,* Wang *et al.* (1998) *Science* 280: 1077-1082; Chee *et al.* (1996) 274: 610-614), various polymerase chain reaction (PCR) assays (*e.g.* that specifically amplify only the mutated variant), various ligase assays, nucleic acid sequencing, and the like. Mutations in polypeptides can easily be detected by changes in polypeptide mobility (*e.g.*
25 electrophoretically), by mass spectroscopy, by the use of antibodies specific to the mutated polypeptide and the like.

III. Mitigation of symptoms of lipodystrophy, obesity, diabetes, atherosclerosis and related pathologies.

[0130] In still another embodiment, this invention provides methods and
30 compositions to mitigate symptoms of lipodystrophy, obesity, diabetes, atherosclerosis

and related pathologies. The methods generally involve increasing or decreasing lipin protein concentration and/or activity in or near a cell, *e.g.*, an adipocyte or an adipocyte precursor. This can be accomplished by transforming a cell (*e.g.* an adipocyte precursor cell) so that it expresses altered (*e.g.* elevated) levels of endogenous *Lpin1/LPIN1* or so that it expresses lipin from an exogenous transfected *Lpin1/LPIN1* nucleic acid, or by contacting the cell(s) with lipin protein(s) through local or systemic administration of a lipin protein, or agent which modulates the expression and/or activity of lipin.

[0131] As used herein, the term "adipocyte precursor cells" refers to any or all of those cells that have the capacity to ultimately form, or contribute to the formation of, new fat tissue. This includes various cells in different stages of differentiation. Adipocyte progenitor cells also include cells that have been isolated and manipulated *in vitro*, *e.g.*, subjected to stimulation with agents such as hormones, cytokines, growth factors or even genetically engineered cells.

[0132] The term "adipocyte precursor cell" is also used to particularly refer to those cells that are located within, are in contact with, or migrate towards (*i.e.*, "home to"), adipocyte progenitor tissue and which cells directly or indirectly stimulate the formation of fat tissue. As such, the progenitor cells may be cells that ultimately differentiate into mature adipocytes themselves, or cells that, upon stimulation, attract further progenitor cells or promote nearby cells to differentiate into adipocytes. Cells affecting adipocyte formation indirectly may do so by the elaboration of various growth factors or cytokines, or by their physical interaction with other cell types.

A) Transformation of cells to increase lipin production.

[0133] In a preferred embodiment, the *Lpin1/LPIN1* nucleic acids (*e.g.*, cDNA(s)) are cloned into gene therapy vectors that are competent to transfect cells (such as human or other mammalian cells) *in vitro* and/or *in vivo*.

[0134] Several approaches for introducing nucleic acids into cells *in vivo*, *ex vivo* and *in vitro* have been used. These include lipid or liposome based gene delivery (WO 96/18372; WO 93/24640; Mannino and Gould-Fogerite (1988) *BioTechniques* 6(7): 682-691; Rose U.S. Pat No. 5,279,833; WO 91/06309; and Felgner *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84: 7413-7414) and replication-defective retroviral vectors harboring a

therapeutic polynucleotide sequence as part of the retroviral genome (see, *e.g.*, Miller *et al.* (1990) *Mol. Cell. Biol.* 10:4239 (1990); Kolberg (1992) *J. NIH Res.* 4: 43, and Cornetta *et al.* (1991) *Hum. Gene Ther.* 2: 215).

[0135] For a review of gene therapy procedures, see, *e.g.*, Anderson, *Science* (1992) 256: 808-813; Nabel and Felgner (1993) *TIBTECH* 11: 211-217; Mitani and Caskey (1993) *TIBTECH* 11: 162-166; Mulligan (1993) *Science*, 926-932; Dillon (1993) *TIBTECH* 11: 167-175; Miller (1992) *Nature* 357: 455-460; Van Brunt (1988) *Biotechnology* 6(10): 1149-1154; Vigne (1995) *Restorative Neurology and Neuroscience* 8: 35-36; Kremer and Perricaudet (1995) *British Medical Bulletin* 51(1) 31-44; Haddada *et al.* (1995) in *Current Topics in Microbiology and Immunology*, Doerfler and Böhm (eds) Springer-Verlag, Heidelberg Germany; and Yu *et al.*, (1994) *Gene Therapy*, 1:13-26.

[0136] Widely used retroviral vectors include those based upon murine leukemia virus (MuLV), gibbon ape leukemia virus (GaLV), Simian Immuno deficiency virus (SIV), human immuno deficiency virus (HIV), and combinations thereof. See, *e.g.*, Buchscher *et al.* (1992) *J. Virol.* 66(5) 2731-2739; Johann *et al.* (1992) *J. Virol.* 66 (5):1635-1640 (1992); Sommerfelt *et al.*, (1990) *Virol.* 176:58-59; Wilson *et al.* (1989) *J. Virol.* 63:2374-2378; Miller *et al.*, *J. Virol.* 65:2220-2224 (1991); Wong-Staal *et al.*, PCT/US94/05700, and Rosenberg and Fauci (1993) in *Fundamental Immunology*, Third Edition Paul (ed) Raven Press, Ltd., New York and the references therein, and Yu *et al.*, *Gene Therapy* (1994) *supra*). The vectors are optionally pseudotyped to extend the host range of the vector to cells which are not infected by the retrovirus corresponding to the vector. The vesicular stomatitis virus envelope glycoprotein (VSV-G) has been used to construct VSV-G-pseudotyped HIV vectors which can infect hematopoietic stem cells (Naldini *et al.* (1996) *Science* 272:263, and Akkina *et al.* (1996) *J Virol* 70:2581).

[0137] Adeno-associated virus (AAV)-based vectors are also used to transduce cells with target nucleic acids, *e.g.*, in the *in vitro* production of nucleic acids and peptides, and in *in vivo* and *ex vivo* gene therapy procedures. See, West *et al.* (1987) *Virology* 160:38-47; Carter *et al.* (1989) U.S. Patent No. 4,797,368; Carter *et al.* WO 93/24641 (1993); Kotin (1994) *Human Gene Therapy* 5:793-801; Muzyczka (1994) *J. Clin. Invest.* 94:1351 for an overview of AAV vectors. Construction of recombinant AAV vectors are described in a number of publications, including Lebkowski, U.S. Pat. No. 5,173,414;

Tratschin *et al.* (1985) *Mol. Cell. Biol.* 5(11):3251-3260; Tratschin, *et al.* (1984) *Mol. Cell. Biol.*, 4: 2072-2081; Hermonat and Muzyczka (1984) *Proc. Natl. Acad. Sci. USA*, 81: 6466-6470; McLaughlin *et al.* (1988) and Samulski *et al.* (1989) *J. Virol.*, 63:03822-3828. Cell lines that can be transformed by rAAV include those described in Lebkowski *et al.*

5 (1988) *Mol. Cell. Biol.*, 8:3988-3996. Other suitable viral vectors include herpes virus and vaccinia virus.

[0138] U.S. Patents 5,942,496 and 5,763,416 disclose methods, compositions, kits and devices for use in transferring nucleic acids into bone cells *in situ* and/or for stimulating bone progenitor cells (*see also*, Evans and Robbins (1995) *J. Bone and Joint*

10 *Surgery*, 77-A(7):1103-1114, Wolff *et al.* (1992) *J. Cell Sci.*, 103:1249-1259).

B) Administration of exogenously produced lipin or inhibitors or lipin expression or activity.

[0139] The lipin proteins (or biologically active fragments thereof) of this invention are or inhibitors of lipin expression and/or activity are useful for intravenous, parenteral, topical, oral, or local administration (*e.g.*, by aerosol or transdermally). Particularly preferred modes of administration include intra-arterial injection, injection into fracture sites or delivery in a biodegradable matrix. The lipin proteins are typically combined with a pharmaceutically acceptable carrier (excipient) to form a pharmacological composition. Pharmaceutically acceptable carriers can contain a

15 physiologically acceptable compound that acts, for example, to stabilize the composition or to increase or decrease the absorption of the agent. Physiologically acceptable compounds can include, for example, carbohydrates, such as glucose, sucrose, or dextrans, antioxidants, such as ascorbic acid or glutathione, chelating agents, low molecular weight proteins, compositions that reduce the clearance or hydrolysis of the anti-mitotic agents, or

20 excipients or other stabilizers and/or buffers.

[0140] Other physiologically acceptable compounds include wetting agents, emulsifying agents, dispersing agents or preservatives that are particularly useful for preventing the growth or action of microorganisms. Various preservatives are well known and include, for example, phenol and ascorbic acid. One skilled in the art would

30 appreciate that the choice of a pharmaceutically acceptable carrier, including a

physiologically acceptable compound depends, for example, on the route of administration of the anti-mitotic agent and on the particular physio-chemical characteristics of the agent.

[0141] The pharmaceutical compositions can be administered in a variety of unit dosage forms depending upon the method of administration. For example, unit dosage forms suitable for oral administration include powder, tablets, pills, capsules and lozenges. It is recognized that the lipin protein(s), if administered orally, are preferably protected from digestion. This is typically accomplished either by complexing the protein with a composition to render it resistant to acidic and enzymatic hydrolysis or by packaging the protein in an appropriately resistant carrier such as a liposome. Means of protecting compounds from digestion are well known in the art (*see, e.g.*, U.S. Patent 5,391,377 describing lipid compositions for oral delivery of therapeutic agents).

[0142] The pharmaceutical compositions of this invention are particularly useful for topical administration *e.g.*, in surgical wounds to treat facilitate bone reconstruction and/or repair. In another embodiment, the compositions are useful for parenteral administration, such as intravenous administration or administration into a body cavity or lumen of an organ. The compositions for administration will commonly comprise a solution of the lipin protein dissolved in a pharmaceutically acceptable carrier, preferably an aqueous carrier for water-soluble proteins. A variety of carriers can be used, *e.g.*, buffered saline and the like. These solutions are sterile and generally free of undesirable matter. These compositions may be sterilized by conventional, well-known sterilization techniques. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, toxicity adjusting agents and the like, for example, sodium acetate, sodium chloride, potassium chloride, calcium chloride, sodium lactate and the like.

[0143] The concentration of lipin protein in these formulations can vary widely, and will be selected primarily based on fluid volumes, viscosities, body weight and the like in accordance with the particular mode of administration selected and the patient's needs. Typically the lipin proteins are utilized in the form of a pharmaceutically acceptable solution (including reconstitution from a lyophilized form). It is optimal to solubilize the protein at concentrations of at least about 1 mg/ml, preferably about 2 to 8 mg/ml, so that a pharmaceutically effective amount of protein can be delivered without

undue volumes of carrier being necessary. For some applications, concentrations above 2 mg/ml may be desirable.

[0144] As alluded to above, the dosage regimen will be determined by the clinical indication being addressed, as well as by various patient variables (e.g. weight, age, sex) and clinical presentation (e.g. extent of injury, site of injury, etc.). In general, the dosage of lipin protein will be in the range of from 1 to about 10000 μ g, preferably from about about 10 to 1000 μ g, more preferably from about 10 to 100 μ g.

C) Inhibition of fat accumulation.

[0145] It is demonstrated herein that lipin deficiency in fld mice is associated with a major reduction in total body fat. In certain embodiments, this invention contemplates the reduction of body fat by inhibiting endogenous lipin expression or activity. Such methods are particularly contemplated in pathologies (e.g. obesity), particularly where lipin is overexpressed.

[0146] Lipin expression can upregulated or inhibited using a wide variety of approaches known to those of skill in the art. For example, methods of inhibiting lipin expression include, but are not limited to antisense molecules, lipin specific ribozymes, lipin specific catalytic DNAs, intrabodies directed against lipin proteins, RNAi, gene therapy approaches that knock out lipins, and small organic molecules that inhibit lipin expression/overexpression or that block receptor activity that is required to induce lipin expression. It will be appreciated that the methods used to alter lipin expression/activity can generally also be used to alter expression/activity of lipin homologues.

1) Antisense approaches.

[0147] Lipin gene expression can be downregulated or entirely inhibited by the use of antisense molecules. An "antisense sequence or antisense nucleic acid" is a nucleic acid that is complementary to the coding lipin mRNA nucleic acid sequence or a subsequence thereof. Binding of the antisense molecule to the lipin mRNA interferes with normal translation of the lipin polypeptide.

[0148] Thus, in accordance with preferred embodiments of this invention, preferred antisense molecules include oligonucleotides and oligonucleotide analogs that

are hybridizable with lipin messenger RNA. This relationship is commonly denominated as "antisense." The oligonucleotides and oligonucleotide analogs are able to inhibit the function of the RNA, either its translation into protein, its translocation into the cytoplasm, or any other activity necessary to its overall biological function. The failure of the messenger RNA to perform all or part of its function results in a reduction or complete inhibition of expression of lipin polypeptides.

[0149] In the context of this invention, the term "oligonucleotide" refers to a polynucleotide formed from naturally-occurring bases and/or cyclofuranosyl groups joined by native phosphodiester bonds. This term effectively refers to naturally-occurring species or synthetic species formed from naturally-occurring subunits or their close homologs. The term "oligonucleotide" may also refer to moieties which function similarly to oligonucleotides, but which have non naturally-occurring portions. Thus, oligonucleotides may have altered sugar moieties or inter-sugar linkages. Exemplary among these are the phosphorothioate and other sulfur containing species that are known for use in the art. In accordance with some preferred embodiments, at least one of the phosphodiester bonds of the oligonucleotide has been substituted with a structure which functions to enhance the ability of the compositions to penetrate into the region of cells where the RNA whose activity is to be modulated is located. It is preferred that such substitutions comprise phosphorothioate bonds, methyl phosphonate bonds, or short chain alkyl or cycloalkyl structures. In accordance with other preferred embodiments, the phosphodiester bonds are substituted with structures which are, at once, substantially non-ionic and non-chiral, or with structures which are chiral and enantiomerically specific. Persons of ordinary skill in the art will be able to select other linkages for use in the practice of the invention.

[0150] In one particularly preferred embodiment, the internucleotide phosphodiester linkage is replaced with a peptide linkage. Such peptide nucleic acids tend to show improved stability, penetrate the cell more easily, and show enhanced affinity for their target. Methods of making peptide nucleic acids are known to those of skill in the art (*see, e.g.*, U.S. Patent Nos: 6,015,887, 6,015,710, 5,986,053, 5,977,296, 5,902,786, 5,864,010, 5,786,461, 5,773,571, 5,766,855, 5,736,336, 5,719,262, and 5,714,331).

[0151] Oligonucleotides may also include species that contain at least some modified base forms. Thus, purines and pyrimidines other than those normally found in nature may be so employed. Similarly, modifications on the furanosyl portions of the nucleotide subunits may also be effected, as long as the essential tenets of this invention are adhered to. Examples of such modifications are 2'-O-alkyl- and 2'-halogen-substituted nucleotides. Some specific examples of modifications at the 2' position of sugar moieties which are useful in the present invention are OH, SH, SCH₃, F, OCH₃, OCN, O(CH₂)[n]NH₂ or O(CH₂)[n]CH₃, where n is from 1 to about 10, and other substituents having similar properties.

[0152] Such oligonucleotides are best described as being functionally interchangeable with natural oligonucleotides or synthesized oligonucleotides along natural lines, but which have one or more differences from natural structure. All such analogs are comprehended by this invention so long as they function effectively to hybridize with messenger RNA of lipin to inhibit the function of that RNA.

[0153] The oligonucleotides in accordance with this invention preferably comprise from about 3 to about 50 subunits. It is more preferred that such oligonucleotides and analogs comprise from about 8 to about 25 subunits and still more preferred to have from about 12 to about 20 subunits. As will be appreciated, a subunit is a base and sugar combination suitably bound to adjacent subunits through phosphodiester or other bonds.

The oligonucleotides used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors, including Applied Biosystems. Any other means for such synthesis may also be employed, however, the actual synthesis of the oligonucleotides is well within the talents of the routineer. It is also well known to prepare other oligonucleotide such as phosphorothioates and alkylated derivatives.

[0154] Using the known sequence of the lipin gene(s)/cDNA(s) identified herein, appropriate and effective antisense oligonucleotide sequences can be readily determined.

2) Catalytic RNAs and DNAs

a) Ribozymes.

[0155] In another approach, lipin expression can be inhibited by the use of ribozymes. As used herein, "ribozymes" include RNA molecules that contain anti-sense sequences for specific recognition, and an RNA-cleaving enzymatic activity. The catalytic strand cleaves a specific site in a target (lipin) RNA, preferably at greater than stoichiometric concentration. Two "types" of ribozymes are particularly useful in this invention, the hammerhead ribozyme (Rossi *et al.* (1991) *Pharmac. Ther.* 50: 245-254) and the hairpin ribozyme (Hampel *et al.* (1990) *Nucl. Acids Res.* 18: 299-304, and U.S. Pat. No. 5,254,678).

[0156] Because both hammerhead and hairpin ribozymes are catalytic molecules having antisense and endoribonucleotidase activity, ribozyme technology has emerged as a powerful extension of the antisense approach to gene inactivation. The ribozymes of the invention typically consist of RNA, but such ribozymes may also be composed of nucleic acid molecules comprising chimeric nucleic acid sequences (such as DNA/RNA sequences) and/or nucleic acid analogs (e.g., phosphorothioates).

[0157] Accordingly, within one aspect of the present invention ribozymes are provided which have the ability to inhibit lipin expression. Such ribozymes can be in the form of a "hammerhead" (for example, as described by Forster and Symons (1987) *Cell* 48: 211-220,; Haseloff and Gerlach (1988) *Nature* 328: 596-600; Walbot and Bruening (1988) *Nature* 334: 196; Haseloff and Gerlach (1988) *Nature* 334: 585) or a "hairpin" (see, e.g. U.S. Patent 5,254,678 and Hampel *et al.*, European Patent Publication No. 0 360 257, published Mar. 26, 1990), and have the ability to specifically target, cleave and lipin nucleic acids.

Sub
25
A1 [0158] The sequence requirement for the hairpin ribozyme is any RNA sequence consisting of NNNBN*GUCNNNNNN (where N*G is the cleavage site, where B is any of G, C, or U, and where N is any of G, U, C, or A) (SEQ ID NO: ____). Suitable lipin of recognition or target sequences for hairpin ribozymes can be readily determined from the lipin sequence(s) identified herein.

[0159] The preferred sequence at the cleavage site for the hammerhead ribozyme is any RNA sequence consisting of NUX (where N is any of G, U, C, or A and X represents C, U, or A) can be targeted. Accordingly, the same target within the hairpin leader sequence, GUC, is useful for the hammerhead ribozyme. The additional
5 nucleotides of the hammerhead ribozyme or hairpin ribozyme is determined by the target flanking nucleotides and the hammerhead consensus sequence (*see* Ruffner *et al.* (1990) *Biochemistry* 29: 10695-10702).

[0160] Cech *et al.* (U.S. Patent 4,987,071,) has disclosed the preparation and use of certain synthetic ribozymes which have endoribonuclease activity. These ribozymes
10 are based on the properties of the Tetrahymena ribosomal RNA self-splicing reaction and require an eight base pair target site. A temperature optimum of 50°C. is reported for the endoribonuclease activity. The fragments that arise from cleavage contain 5' phosphate and 3' hydroxyl groups and a free guanosine nucleotide added to the 5' end of the cleaved RNA. The preferred ribozymes of this invention hybridize efficiently to target sequences
15 at physiological temperatures, making them particularly well suited for use *in vivo*.

[0161] The ribozymes of this invention, as well as DNA encoding such ribozymes and other suitable nucleic acid molecules can be chemically synthesized using methods well known in the art for the synthesis of nucleic acid molecules. Alternatively, Promega, Madison, Wis., USA, provides a series of protocols suitable for the production of RNA
20 molecules such as ribozymes. The ribozymes also can be prepared from a DNA molecule or other nucleic acid molecule (which, upon transcription, yields an RNA molecule) operably linked to an RNA polymerase promoter, *e.g.*, the promoter for T7 RNA polymerase or SP6 RNA polymerase. Such a construct may be referred to as a vector. Accordingly, also provided by this invention are nucleic acid molecules, *e.g.*, DNA or
25 cDNA, coding for the ribozymes of this invention. When the vector also contains an RNA polymerase promoter operably linked to the DNA molecule, the ribozyme can be produced *in vitro* upon incubation with the RNA polymerase and appropriate nucleotides. In a separate embodiment, the DNA may be inserted into an expression cassette (*see, e.g.*, Cotten and Birnstiel (1989) *EMBO J* 8(12):3861-3866; Hempel *et al.* (1989) *Biochem.* 28:
30 4929-4933, *etc.*).

[0162] After synthesis, the ribozyme can be modified by ligation to a DNA molecule having the ability to stabilize the ribozyme and make it resistant to RNase. Alternatively, the ribozyme can be modified to the phosphothio analog for use in liposome delivery systems. This modification also renders the ribozyme resistant to endonuclease activity.

[0163] The ribozyme molecule also can be in a host prokaryotic or eukaryotic cell in culture or in the cells of an organism/patient. Appropriate prokaryotic and eukaryotic cells can be transfected with an appropriate transfer vector containing the DNA molecule encoding a ribozyme of this invention. Alternatively, the ribozyme molecule, including nucleic acid molecules encoding the ribozyme, may be introduced into the host cell using traditional methods such as transformation using calcium phosphate precipitation (Dubensky *et al.* (1984) *Proc. Natl. Acad. Sci., USA*, 81: 7529-7533), direct microinjection of such nucleic acid molecules into intact target cells (Acsadi *et al.* (1991) *Nature* 352: 815-818), and electroporation whereby cells suspended in a conducting solution are subjected to an intense electric field in order to transiently polarize the membrane, allowing entry of the nucleic acid molecules. Other procedures include the use of nucleic acid molecules linked to an inactive adenovirus (Cotton *et al.* (1990) *Proc. Natl. Acad. Sci., USA*, 89 :6094), lipofection (Felgner *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 84: 7413-7417), microprojectile bombardment (Williams *et al.* (1991) *Proc. Natl. Acad. Sci., USA*, 88: 2726-2730), polycation compounds such as polylysine, receptor specific ligands, liposomes entrapping the nucleic acid molecules, spheroplast fusion whereby *E coli* containing the nucleic acid molecules are stripped of their outer cell walls and fused to animal cells using polyethylene glycol, viral transduction, (Cline *et al.*, (1985) *Pharmac. Ther.* 29: 69; and Friedmann *et al.* (1989) *Science* 244: 1275), and DNA ligand (Wu *et al* (1989) *J. Biol. Chem.* 264: 16985-16987), as well as psoralen inactivated viruses such as Sendai or Adenovirus. In one preferred embodiment, the ribozyme is introduced into the host cell utilizing a lipid, a liposome or a retroviral vector.

[0164] When the DNA molecule is operatively linked to a promoter for RNA transcription, the RNA can be produced in the host cell when the host cell is grown under suitable conditions favoring transcription of the DNA molecule. The vector can be, but is not limited to, a plasmid, a virus, a retrotransposon or a cosmid. Examples of such vectors are disclosed in U.S. Pat. No. 5,166,320. Other representative vectors include, but are not

limited to adenoviral vectors (e.g., WO 94/26914, WO 93/9191; Kolls *et al.* (1994) PNAS 91(1):215-219; Kass-Eisler *et al.*, (1993) *Proc. Natl. Acad. Sci., USA*, 90(24): 11498-502, Guzman *et al.* (1993) *Circulation* 88(6): 2838-48, 1993; Guzman *et al.* (1993) *Cir. Res.* 73(6):1202-1207, 1993; Zabner *et al.* (1993) *Cell* 75(2): 207-216; Li *et al.* (1993) *Hum Gene Ther.* 4(4): 403-409; Caillaud *et al.* (1993) *Eur. J Neurosci.* 5(10): 1287-1291), adeno-associated vector type 1 ("AAV-1") or adeno-associated vector type 2 ("AAV-2") (see WO 95/13365; Flotte *et al.* (1993) *Proc. Natl. Acad. Sci., USA*, 90(22):10613-10617), retroviral vectors (e.g., EP 0 415 731; WO 90/07936; WO 91/02805; WO 94/03622; WO 93/25698; WO 93/25234; U.S. Pat. No. 5,219,740; WO 93/11230; WO 93/10218) and herpes viral vectors (e.g., U.S. Pat. No. 5,288,641). Methods of utilizing such vectors in gene therapy are well known in the art, see, for example, Larrick and Burck (1991) *Gene Therapy: Application of Molecular Biology*, Elsevier Science Publishing Co., Inc., New York, New York, and Kreigler (1990) *Gene Transfer and Expression: A Laboratory Manual*, W.H. Freeman and Company, New York.

[0165] To produce ribozymes *in vivo* utilizing vectors, the nucleotide sequences coding for ribozymes are preferably placed under the control of a strong promoter such as the lac, SV40 late, SV40 early, or lambda promoters. Ribozymes are then produced directly from the transfer vector *in vivo*. Suitable transfactor vectors for *in vivo* expression are discussed below.

20 **b) Catalytic DNA**

[0166] In a manner analogous to ribozymes, DNAs are also capable of demonstrating catalytic (e.g. nuclease) activity. While no such naturally-occurring DNAs are known, highly catalytic species have been developed by directed evolution and selection. Beginning with a population of 10^{14} DNAs containing 50 random nucleotides, successive rounds of selective amplification, enriched for individuals that best promote the Pb^{2+} -dependent cleavage of a target ribonucleoside 3'-O-P bond embedded within an otherwise all-DNA sequence. By the fifth round, the population as a whole carried out this reaction at a rate of 0.2 min^{-1} . Based on the sequence of 20 individuals isolated from this population, a simplified version of the catalytic domain that operates in an intermolecular context with a turnover rate of 1 min^{-1} (see, e.g., Breaker and Joyce (1994) *Chem Biol* 4: 223-229).



[0167] In later work, using a similar strategy, a DNA enzyme was made that could cleave almost any targeted RNA substrate under simulated physiological conditions. The enzyme is comprised of a catalytic domain of 15 deoxynucleotides, flanked by two substrate-recognition domains of seven to eight deoxynucleotides each. The RNA substrate is bound through Watson-Crick base pairing and is cleaved at a particular phosphodiester located between an unpaired purine and a paired pyrimidine residue. Despite its small size, the DNA enzyme has a catalytic efficiency (k_{cat}/K_m) of approximately $10^9 \text{ M}^{-1} \text{ min}^{-1}$ under multiple turnover conditions, exceeding that of any other known nucleic acid enzyme. By changing the sequence of the substrate-recognition domains, the DNA enzyme can be made to target different RNA substrates (Santoro and Joyce (1997) *Proc. Natl. Acad. Sci., USA*, 94(9): 4262-4266). Modifying the appropriate targeting sequences (*e.g.* as described by Santoro and Joyce, *supra.*) the DNA enzyme can easily be retargeted to lipin mRNA thereby acting like a ribozyme.

3) Knocking out lipin

[0168] In another approach, lipin can be inhibited/downregulated simply by "knocking out" the gene.

[0169] In certain embodiments, this invention provides animals in which the lipin gene is knocked out". Such animals can be heterozygous or homozygous for the knockout.

[0170] Typically this is accomplished by disrupting the lipin gene(s), the promoter regulating the lipin gene(s) or sequences between the endogenous promoter(s) and the gene(s). Such disruption can be specifically directed to lipin nucleic acids (*e.g.* lipin1, and/or lipin2, and/or lipin3) by homologous recombination where a "knockout construct" contains flanking sequences complementary to the domain to which the construct is targeted. Insertion of the knockout construct (*e.g.* into a lipin gene) results in disruption of that gene.

[0171] The phrases "disruption of the gene" and "gene disruption" refer to insertion of a nucleic acid sequence into one region of the native DNA sequence (usually one or more exons) and/or the promoter region of a gene so as to decrease or prevent expression of that gene in the cell as compared to the wild-type or naturally occurring

sequence of the gene. By way of example, a nucleic acid construct can be prepared containing a DNA sequence encoding an antibiotic resistance gene which is inserted into the DNA sequence that is complementary to the DNA sequence (promoter and/or coding region) to be disrupted. When this nucleic acid construct is then transfected into a cell, the construct will integrate into the genomic DNA. Thus, the cell and its progeny will no longer express the gene or will express it at a decreased level, as the DNA is now disrupted by the antibiotic resistance gene.

[0172] Knockout constructs can be produced by standard methods known to those of skill in the art. The knockout construct can be chemically synthesized or assembled, *e.g.*, using recombinant DNA methods. The DNA sequence to be used in producing the knockout construct is digested with a particular restriction enzyme selected to cut at a location(s) such that a new DNA sequence encoding a marker gene can be inserted in the proper position within this DNA sequence. The proper position for marker gene insertion is that which will serve to prevent expression of the native lipin gene; this position will depend on various factors such as the restriction sites in the sequence to be cut, and whether an exon sequence or a promoter sequence, or both is (are) to be interrupted (*i.e.*, the precise location of insertion necessary to inhibit promoter function or to inhibit synthesis of the native exon). Preferably, the enzyme selected for cutting the DNA will generate a longer arm and a shorter arm, where the shorter arm is at least about 300 base pairs (bp). In some cases, it will be desirable to actually remove a portion or even all of one or more exons of the gene to be suppressed so as to keep the length of the knockout construct comparable to the original genomic sequence when the marker gene is inserted in the knockout construct. In these cases, the genomic DNA is cut with appropriate restriction endonucleases such that a fragment of the proper size can be removed.

[0173] The marker gene can be any nucleic acid sequence that is detectable and/or assayable, however typically it is an antibiotic resistance gene or other gene whose expression or presence in the genome can easily be detected. The marker gene is usually operably linked to its own promoter or to another strong promoter from any source that will be active or can easily be activated in the cell into which it is inserted; however, the marker gene need not have its own promoter attached as it may be transcribed using the promoter of the gene to be suppressed. In addition, the marker gene will normally have a polyA sequence attached to the 3' end of the gene; this sequence serves to terminate

transcription of the gene. Preferred marker genes are any antibiotic resistance gene including, but not limited to *neo* (the neomycin resistance gene) and *beta-gal* (beta-galactosidase).

[0174] After the genomic DNA sequence has been digested with the appropriate restriction enzymes, the marker gene sequence is ligated into the genomic DNA sequence using methods well known to the skilled artisan (*see, e.g.,* Berger and Kimmel, *Guide to Molecular Cloning Techniques, Methods in Enzymology* volume 152 Academic Press, Inc., San Diego, CA; Sambrook *et al.* (1989) *Molecular Cloning - A Laboratory Manual* (2nd ed.) Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor Press, NY; and *Current Protocols in Molecular Biology*, F.M. Ausubel *et al.*, eds., Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (1994) Supplement). The ends of the DNA fragments to be ligated are rendered compatible, *e.g.,* by either cutting the fragments with enzymes that generate compatible ends, or by blunting the ends prior to ligation. Blunting is done using methods well known in the art, such as for example by the use of Klenow fragment (DNA polymerase I) to fill in sticky ends.

[0175] The production of knockout constructs and their use to produce knockout mice is well known to those of skill in the art (*see, e.g.,* Dorfman *et al.* (1996) *Oncogene* 13: 925-931). The knockout constructs can be delivered to cells *in vivo* using gene therapy delivery vehicles (*e.g.* retroviruses, liposomes, lipids, dendrimers, *etc.*) as described above. Methods of knocking out genes are well described in the literature and essentially routine to those of skill in the art (*see, e.g.,* Thomas *et al.* (1986) *Cell* 44(3): 419-428; Thomas, *et al.* (1987) *Cell* 51(3): 503-512]; Jasin and Berg (1988) *Genes & Development* 2: 1353-1363; Mansour, *et al.* (1988) *Nature* 336: 348-352; Brinster, *et al.* (1989) *Proc Natl Acad Sci* 86: 7087-7091; Capecchi (1989) *Trends in Genetics* 5(3): 70-76; Frohman and Martin (1989) *Cell* 56: 145-147; Hasty, *et al.* (1991) *Mol Cell Biol* 11(11): 5586-5591; Jeannotte, *et al.* (1991) *Mol Cell Biol*. 11(11): 5578-5585; and Mortensen, *et al.* (1992) *Mol Cell Biol*. 12(5): 2391-2395.

[0176] The use of homologous recombination to alter expression of endogenous genes is also described in detail in U.S. Patent 5,272,071, WO 91/09955, WO 93/09222, WO 96/29411, WO 95/31560, and WO 91/12650.

[0177] Production of the knockout animals of this invention is not dependent on the availability of ES cells. In various embodiments, knockout animals of this invention can be produced using methods of somatic cell nuclear transfer. In preferred embodiments using such an approach, a somatic cell is obtained from the species in which the lipin gene is to be knocked out. The cell is transfected with a construct that introduces a disruption in the lipin gene (*e.g.* via heterologous recombination) as described herein. Cells harboring a knocked out lipin gene are selected as described herein. The nucleus of such cells harboring the knockout is then placed in an unfertilized enucleated egg (*e.g.*, eggs from which the natural nuclei have been removed by microsurgery). Once the transfer is complete, the recipient eggs contained a complete set of genes, just as they would if they had been fertilized by sperm. The eggs are then cultured for a period before being implanted into a host mammal (of the same species that provided the egg) where they are carried to term, culminating in the berth of a transgenic animal comprising a nucleic acid construct containing one or more disrupted lipin genes.

[0178] The production of viable cloned mammals following nuclear transfer of cultured somatic cells has been reported for a wide variety of species including, but not limited to frogs (McKinnell (1962) *J. Hered.* 53, 199–207), calves (Kato *et al.* (1998) *Science* 262: 2095–2098), sheep (Campbell *et al.* (1996) *Nature* 380: 64–66), mice (Wakayama and Yanagimachi (1999) *Nat. Genet.* 22: 127–128), goats (Baguisi *et al.* (1999) *Nat. Biotechnol.* 17: 456–461), monkeys (Meng *et al.* (1997) *Biol. Reprod.* 57: 454–459), and pigs (Bishop *et al.* (2000) *Nature Biotechnology* 18: 1055–1059). Nuclear transfer methods have also been used to produce clones of transgenic animals. Thus, for example, the production of transgenic goats carrying the human antithrobin III gene by somatic cell nuclear transfer has been reported (Baguisi *et al.* (1999) *Nature Biotechnology* 17: 456–461).

[0179] Using methods of nuclear transfer as described in these and other references, cell nuclei derived from differentiated fetal or adult, mammalian cells are transplanted into enucleated mammalian oocytes of the same species as the donor nuclei. The nuclei are reprogrammed to direct the development of cloned embryos, which can then be transferred into recipient females to produce fetuses and offspring, or used to produce cultured inner cell mass (CICM) cells. The cloned embryos can also be combined with fertilized embryos to produce chimeric embryos, fetuses and/or offspring.

[0180] Somatic cell nuclear transfer also allows simplification of transgenic procedures by working with a differentiated cell source that can be clonally propagated. This eliminates the need to maintain the cells in an undifferentiated state, thus, genetic modifications, both random integration and gene targeting, are more easily accomplished.

5 Also by combining nuclear transfer with the ability to modify and select for these cells *in vitro*, this procedure is more efficient than previous transgenic embryo techniques.

[0181] Nuclear transfer techniques or nuclear transplantation techniques are known in the literature. See, in particular, Campbell *et al.* (1995) *Theriogenology*, 43:181; Collas *et al.* (1994) *Mol. Report Dev.*, 38:264-267; Keefer *et al.* (1994) *Biol. Reprod.*, 50:935-939; Sims *et al.* (1993) *Proc. Natl. Acad. Sci., USA*, 90:6143-6147; WO 94/26884; 10 WO 94/24274, WO 90/03432, U.S. Patents 5,945,577, 4,944,384, 5,057,420 and the like.

4) Intrabodies.

[0182] In still another embodiment, lipin expression/activity is inhibited by transfecting the subject cell(s) (*e.g.*, cells of the vascular endothelium) with a nucleic acid construct that expresses an intrabody. An intrabody is an intracellular antibody, in this 15 case, capable of recognizing and binding to a lipin polypeptide. The intrabody is expressed by an "antibody cassette", containing a sufficient number of nucleotides coding for the portion of an antibody capable of binding to the target (lipin polypeptide) operably linked to a promoter that will permit expression of the antibody in the cell(s) of interest.

20 The construct encoding the intrabody is delivered to the cell where the antibody is expressed intracellularly and binds to the target lipin, thereby disrupting the target from its normal action. This antibody is sometimes referred to as an "intrabody".

[0183] In one preferred embodiment, the "intrabody gene" (antibody) of the antibody cassette would utilize a cDNA, encoding heavy chain variable (V_H) and light chain variable (V_L) domains of an antibody which can be connected at the DNA level by 25 an appropriate oligonucleotide as a bridge of the two variable domains, which on translation, form a single peptide (referred to as a single chain variable fragment, "sFv") capable of binding to a target such as a lipin protein. The intrabody gene preferably does not encode an operable secretory sequence and thus the expressed antibody remains within 30 the cell.



[0184] Anti-lipin antibodies suitable for use/expression as intrabodies in the methods of this invention can be readily produced by a variety of methods. Such methods include, but are not limited to, traditional methods of raising "whole" polyclonal antibodies, which can be modified to form single chain antibodies, or screening of, *e.g.* phage display libraries to select for antibodies showing high specificity and/or avidity for lipin. Such screening methods are described above in some detail.

[0185] The antibody cassette is delivered to the cell by any of the known means. This discloses the use of a fusion protein comprising a target moiety and a binding moiety. The target moiety brings the vector to the cell, while the binding moiety carries the antibody cassette. Other methods include, for example, Miller (1992) *Nature* 357: 455-460; Anderson (1992) *Science* 256: 808-813; Wu, *et al.* (1988) *J. Biol. Chem.* 263: 14621-14624. For example, a cassette containing these (anti-lipin) antibody genes, such as the sFv gene, can be targeted to a particular cell by a number of techniques including, but not limited to the use of tissue-specific promoters, the use of tissue specific vectors, and the like. Methods of making and using intrabodies are described in detail in U.S. Patent 6,004,940.

5) Small organic molecules.

[0186] In still another embodiment, lipin expression and/or lipin protein activity can be inhibited by the use of small organic molecules. Such molecules include, but are not limited to molecules that specifically bind to the DNA comprising the lipin promoter and/or coding region, molecules that bind to and complex with lipin mRNA, molecules that inhibit the signaling pathway that results in lipin upregulation, and molecules that bind to and/or compete with lipin polypeptides. Small organic molecules effective at inhibiting lipin expression can be identified with routine screening using the methods described herein.

[0187] The methods of inhibiting lipin expression described above are meant to be illustrative and not limiting. In view of the teachings provided herein, other methods of inhibiting lipin will be known to those of skill in the art.

V. Kits.

[0188] In still another embodiment, this invention provides kits for practice of the assays or use of the compositions described herein. In one preferred embodiment, the kits comprise one or more containers containing antibodies and/or nucleic acid probes and/or substrates suitable for detection of *Lpin1* expression and/or activity levels. The kits may optionally include any reagents and/or apparatus to facilitate practice of the assays described herein. Such reagents include, but are not limited to buffers, labels, labeled or unlabeled antibodies, labeled nucleic acids, filter sets for visualization of fluorescent labels, blotting membranes, and the like.

[0189] In another embodiment, the kits can comprise a container containing an lipin protein, or a vector encoding a lipin protein and/or a cell comprising a vector encoding a lipin protein.

[0190] In addition, the kits may include instructional materials containing directions (*i.e.*, protocols) for the practice of the assay methods of this invention or the administration of the compositions described here along with counterindications. While the instructional materials typically comprise written or printed materials they are not limited to such. Any medium capable of storing such instructions and communicating them to an end user is contemplated by this invention. Such media include, but are not limited to electronic storage media (e.g., magnetic discs, tapes, cartridges, chips), optical media (e.g., CD ROM), and the like. Such media may include addresses to internet sites that provide such instructional materials.

EXAMPLES

[0191] The following examples are offered to illustrate, but not to limit the claimed invention.

Example 1**Lipodystrophy in the *fld* Mouse Results From Mutation in a Novel Nuclear Protein**

[0192] Mice carrying mutations in the fatty liver dystrophy (*fld*) gene exhibit features of human lipodystrophy, a genetically heterogeneous group of disorders characterized by loss of body fat, fatty liver, hypertriglyceridemia, insulin resistance, and

increased susceptibility to atherosclerosis (Seip and Trygstad (1996) *Acta Paediatr. Scand. Suppl.* 413: 2-28; Senior and Gellis (1964) *Pediatrics* 33: 593-612; Dörfler *et al.* (1993) *Clin. Investig.* 71: 264-269; Garg (2000) *J. Clin. Endocrinol. Metab.* 85: 1776-1782).

Through positional cloning, we have isolated the responsible gene, and characterized two independent mutant alleles, *fld* and *fld*^{2J}. The gene encodes a novel nuclear protein which we have named lipin (gene symbol *Lpin1*). Consistent with the observed reduction of adipose tissue mass in *fld* and *fld*^{2J} mice, wild-type lipin mRNA is expressed at high levels in adipose tissue and is induced during differentiation of 3T3-L1 preadipocytes. Our results indicate that lipin is required for normal adipose tissue development, and provide a candidate gene for human lipodystrophy. Lipin defines a novel family of nuclear proteins containing at least three members in mammalian species, and homologs in distantly related organisms from man to yeast.

[0193] The initial description of the BALB/cByJ-*fld* mutant mouse (hereafter known as *fld*) focused on two features for which the mutation was named—presence of a triglyceride-filled fatty liver and a progressive neuropathy affecting peripheral nerve (Langner *et al.* (1989) *J. Biol. Chem.* 264: 7994-8003; Langner *et al.* (1991) *J. Biol. Chem.* 266: 11955-11964; Rehnmark *et al.* (1998) *J. Lipid Res.* 39: 2209-2217; Klingenspor *et al.* (1999) *J. Biol. Chem.* 274: 23078-23084). Subsequently, an independent mutant strain, C3H/HeJ-*fld*^{2J} (known as *fld*^{2J}), with the same phenotype and a mutation allelic to *fld* was isolated (Mouse Genome Informatics Project: The Jackson Laboratory: Bar Harbor: Maine. (URL:<http://www.informatics.jax.org>) (1999)). Our further characterization indicates that in addition to fatty liver and neuropathy, both *fld* and *fld*^{2J} mutants exhibit markedly diminished adipose tissue depots with 50-90% reductions in white and brown fat pad mass (Fig. 1a and data not shown). Adipocytes in the affected tissue appear immature, with sparse lipid droplets (compare Fig. 1b and c). Similar reductions in adipose tissue mass and cellular lipid content have been observed in transgenic mice expressing a dominant negative transcription factor that interferes with adipocyte differentiation (Shimomura *et al.* (1998) *Genes. Dev.* 12: 3182-3194). In these mice, it has been shown that reduction in leptin levels that results from diminished adipose tissue mass is responsible for the development of fatty liver, hypertriglyceridemia and insulin resistance (Shimomura *et al.* (1999) *Science* 401: 73-76). Analogously, *fld* mice exhibit significantly reduced plasma leptin levels (Fig. 1d) and insulin resistance, suggesting that

the primary defect in *fld* and *fld*^{2J} mice is impaired adipose tissue development, with other phenotypic features (*i.e.*, fatty liver, hypertriglyceridemia, insulin resistance) occurring as secondary manifestations of the mutation.

[0194] To isolate the gene responsible for the *fld* phenotype we employed a positional cloning strategy. We previously identified three genes in the *fld* critical region (Péterfy *et al.* (1999) *Genomics* 62: 436-444), and now report that one of these, *Lpin1* (formerly *Kiaa0188*), is mutated in both the *fld* and *fld*^{2J} strains.

[0195] Characterization of *Lpin1* revealed a complex rearrangement in the genome of *fld* mice compared to their wild-type littermates. Southern blot analysis using exonic DNA probes from the 5' and 3' ends of the gene indicated the presence of a deletion and duplication, respectively (Fig. 2a). PCR scanning of the gene uncovered additional alterations in *fld* DNA, with primer sets flanking the deleted and duplicated regions failing to amplify *fld* DNA (Fig. 2b). However, when forward or reverse primers from opposite ends of the gene were paired, PCR products were generated from *fld* DNA, indicating an internal inversion. Based on the integration of these results and additional Southern blot, PCR amplification and DNA sequence data (not shown), we determined that the *fld* mutant allele contains three gross abnormalities, depicted schematically in Fig. 2c: (i) the deletion of ~2 kb resulting in the elimination of exons 2 and 3, including the translation initiation site; (ii) the inversion of a large genomic region covering more than 40 kb and containing the majority of the coding sequence; and (iii) duplication of a 0.5-kb segment of the 3' UTR, with one copy at the position it occurs in the wild-type allele and a second copy carried along with the inverted region. The 5' and 3' ends of the *fld* allele remain intact (Fig. 2c and data not shown), indicating that the mutation is confined to *Lpin1*. The molecular mechanisms leading to the *fld* allele are unknown, although, the coincidence of chromosomal breakpoints at the deletion/inversion and inversion/duplication junctions suggests a concerted event rather than a series of independent steps (Okano *et al.* (1991) *J. Neurochem.* 56: 560-567). Furthermore, since no homologous sequences could be identified at the breakpoint regions, a mechanism involving non-homologous recombination is suggested.

[0196] In contrast to *fld*, PCR scanning of genomic DNA from the *fld*^{2J} strain revealed no gross abnormalities in *Lpin1* (not shown). We therefore sequenced lipin

cDNA and genomic DNA prepared from tissues of *fld^{2J}* and wild-type littermates, and detected a single point mutation resulting in a Gly84Arg substitution (Fig. 2d). As shown in Fig. 4d, Gly84 is invariably conserved in homologous sequences from diverse organisms including human, *Drosophila melanogaster*, *Caenorhabditis elegans*,
 5 *Arabidopsis thaliana*, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe* and *Plasmodium falciparum*. These data strongly suggest that the nonconservative mutation in the *fld^{2J}* allele has a functional consequence on lipin activity.

[0197] Northern blot analysis of total RNA from wild-type mouse tissues revealed that lipin mRNA is prominently expressed in adipose tissue, skeletal muscle and testis
 10 (Fig. 3a). Lower level expression was also detected in kidney, lung, brain, and liver (the latter was detectable only with poly A+ RNA). In addition to the ~5-kb transcript observed in these tissues, testis also exhibited an abundant ~3-kb transcript. Aberrant lipin mRNA levels were evident in tissue from mice homozygous for the *fld* and *fld^{2J}* mutations. No mRNA was detectable in *fld* adipose tissue, whereas elevated mRNA levels were
 15 detected in *fld^{2J}* adipose (Fig. 3b). Increased expression in *fld^{2J}* tissue may be the consequence of transcriptional feedback regulation of the *Lpin1* gene due to impaired lipin function. Consistent with impaired adipocyte development in *fld* mice, lipin mRNA expression was induced upon differentiation of 3T3-L1 preadipocytes (Fig. 3c) co-incident with expression of adipsin, a marker of mature adipocytes (Ailhaud *et al.* (1992) *Ann. Rev.*
 20 *Nutr.* 12: 207-233).

[0198] *Lpin1* encodes a novel gene product of 891 amino acids with no similarity to previously characterized proteins or protein domains, although motif searches revealed the presence of a putative nuclear localization signal (NLS) (Fig. 4a). Subcellular
 25 localization of a lipin-GFP fusion protein confirmed that lipin is a nuclear protein (Fig. 4b).

[0199] Database searches identified several mouse and human EST and genomic sequences with significant similarities to *Lpin1*. Using RACE cloning and gene prediction, we identified two *Lpin1*-related mouse genes, *Lpin2* and *Lpin3*, and three human homologs indicating that *Lpins* form a gene family in both organisms (Fig. 4a).
 30 The mouse and human genes form three orthologous pairs as determined by sequence comparison and genomic mapping (Fig. 4c). Multiple sequence alignment of lipin-related

proteins revealed two strongly conserved regions, which we designate N-terminal and C-terminal lipin (NLIP and CLIP) domains (Fig. 4*a* and *d*). In addition to mouse and human proteins, we recognized several predicted proteins with NLIP and CLIP domains from a broad range of eukaryotic organisms (Fig. 4*a*), indicating that these protein domains may be required for the function of lipin and its homologs.

[0200] The identification of lipin has revealed a new factor required for normal adipose tissue development and metabolism. Elucidation of the molecular function of lipin will likely lead to new insights into these processes. The existence of at least two additional *Lpin1*-related genes suggests that members of this novel protein family may have roles in diverse tissues and cellular processes. The human ortholog of the *Lpin1* gene, *LPIN1*, is a potential candidate gene for lipodystrophy, a heterogeneous group of disorders with unknown genetic determinants, except for the *LMNA* gene responsible for Dunnigan-type familial partial lipodystrophy (Cao and Hegele (2000) *Hum. Mol. Genet.* 9: 109-112; Shackleton *et al.* (2000) *Nature Genet.* 24: 153-156). Studies showing linkage of the chromosome 2p21 region harboring *LPIN1* to fat mass (Comuzzie *et al.* (1997) *Nature Genet.* 15: 273-275) and serum leptin levels (Comuzzie *et al.* (1997) *Nature Genet.* 15: 273-275; Rotimi *et al.* (1999) *Diabetes* 48: 643-644; Hager *et al.* (1998) *Nature Genet.* 20: 304-308) support this possibility.

Methods

Mice.

[0201] BALB/cByJ-*+fld* and C3H/HeJ-*+fld^{2J}* mice were obtained from the Jackson Laboratory and bred to produce the phenotypically mutant (*fld/fld*, *fld^{2J}/fld^{2J}*) and wild-type (*+/+* and *+fld*, *+fld^{2J}*) mice used in these studies. Mice were fed a standard laboratory chow diet (Purina 5001) and maintained in a 14:10-hr light-dark cycle. All animal experiments were performed according to guidelines established in the "Guide for the Care and Use of Laboratory Animals".

Adipose tissue and leptin analysis.

[0202] Adipose tissue morphology was assessed in 4 μ m sections of epididymal fat pads stained with hematoxylin and eosin. For leptin measurements, blood was drawn

from the retro-orbital sinus and plasma prepared by centrifugation. Plasma leptin levels were determined with a mouse leptin RIA kit (Linco).

Southern blots.

Sub A2
5 [0203] Mouse genomic DNA was prepared using DNAzol (Gibco/BRL). Blots containing restriction digested DNA (5 μ g/lane) were hybridized to a 257-bp 5' probe corresponding to exons 2 and 3 derived from exon trapping (Péterfy *et al.* (1999) *Genomics* 62: 436-444), or a 225-bp 3' probe corresponding to part of the 3' UTR and generated by PCR (5'-TAC GCA GGG ACA CAT TTC CA-3', SEQ ID NO:_) and 5'-GAG AGA TGC AGC TGC GTC A-3', SEQ ID NO:_). Hybridizations were performed at 65°C in 0.5 M sodium phosphate, pH 7.0, 7% SDS, 1% BSA, and washed at 65°C to a final stringency of 0.1X SSC/ 0.1% SDS. Hybridization signals were detected by phosphorimaging.

Northern blots.

15 [0204] Total RNA was prepared from flash-frozen mouse tissues and 3T3-L1 cells using TRIzol (Gibco/BRL) and 20 μ g per lane was run in 1.2% agarose/formaldehyde gels. Poly A+ RNA from liver was prepared using PolyAtract reagents (Promega) and 2 μ g were analyzed. Hybridizations and washes were performed as for Southern blots, except that the temperature was 63°C. The probe used for Northern blots contained sequence corresponding to nts 1-1264 of lipin cDNA (AF180471). Adipsin cDNA was 20 kindly provided by Dr. B. Spiegelman (Spiegelman *et al.* (1983) *J. Biol. Chem.* 258: 10083-10089).

PCR, RACE and DNA sequencing.

Sub A3
25 [0205] PCR amplification of genomic DNA was performed in an MJ Research PTC-200 thermocycler (1 min 94°C, 45 sec 55°C, 1-2 min 72°C) for 30-32 cycles. Primers for amplification of the inversion breakpoints (Fig. 2b) were: p1, 5'-CCC TTG AGC ACG TTC ACA-3' (SEQ ID NO:_); p2, 5'-CTG ATC GTT GTC AGT CTC T-3' (SEQ ID NO:_); p3, 5'-GGT TGT GGG GAC CCT GGA-3' (SEQ ID NO:_); p4, 5'-GCC TGC TGC AGA TGC GTT-3' (SEQ ID NO:_). RACE cloning of full length cDNAs for *Lpin2* and *Lpin3* was performed using liver cDNA template prepared with the Marathon cDNA

Amplification Kit (Clontech). PCR products were TA-cloned into pCR2.1 (Invitrogen), and sequenced using the Amplicycle sequencing kit (Perkin Elmer) and an ABI model 373A sequencer.

Nuclear localization studies.

Sub
A5
[0206] The entire coding region of the lipin cDNA was amplified from liver cDNA using the primers 5'-GCT CGA ATT CAG ACA ATG AAT TAC GTG GGG CAG CT-3' (SEQ ID NO:_) and 5'-CGT GCA GTC GAC GCT GAG GCT GAA TGC ATG TCC TGG T-3' (SEQ ID NO:_) and cloned as an *EcoRI/SalI* fragment into the pEGFP-N1 vector (Clontech). 3T3-L1 cells were transfected using Lipofectin (Gibco/BRL). 48 hours after transfection, cells were fixed with 4% paraformaldehyde in PBS, stained with Hoechst-33258 dye, and observed with a Zeiss Axiophot fluorescence microscope.

Cell culture.

[0207] The 3T3-L1 preadipocyte cell line (ATCC CL-173) was maintained in DMEM containing 10% fetal bovine serum (basal medium). To induce adipocyte differentiation, cells were grown to confluence (day 0) and then cultured for 3 days in differentiation medium consisting of basal medium plus insulin (10 μ g/ml), dexamethasone (2 μ g/ml), and methylisobutylxanthine (0.5 mM) (Student *et al.* (1980) *J. Biol. Chem.* 255: 4745-4750). On days 4-7, cells were maintained in basal medium with insulin.

Radiation hybrid mapping.

20
Sub
A5
[0208] *Lpin2* and *Lpin3* were mapped using a mouse-hamster radiation hybrid panel (Research Genetics) (McCarthy *et al.* (1997) *Genome Res.* 7: 1153-1161). Oligonucleotide primer pairs derived from the 3' UTR of each gene were as follows: *Lpin2* (5'-GGC GAG ACC CAA TCC CTG A-3', SEQ ID NO:_) and 5'-GGG TCT TCC TCT GTA AGA-3', SEQ ID NO:_) ; *Lpin3* (5'-CCT GGC TTG AGC TTG CCT T-3', SEQ ID NO:_, and 5'-CCC ACG GCA TGC ATC TTC T-3', SEQ ID NO:_.).

GenBank accession numbers.

[0209] *Lpin1*, AF180470; *Lpin2*, AF286723; *Lpin3*, AF286724

Example 2

Further Evidence of the Role of Lipin in Adipogenesis

[0210] Several lines of evidence indicate that lipin is required for normal
5 adipogenesis. These include whole animal studies and studies in isolated cells.

[0211] Lipin deficiency in fld mice is associated with a major reduction in total
body fat (3-4% as compared to 18% in wild-type mice of the same strain), as well as
diminished fat pad mass (10% of normal) for epididymal, inguinal (subcutaneous), and
interscapular brown fat pads. What little adipose tissue is present in fld adipose tissue
10 depots appears abnormal, with very little lipid accumulation in cells and aberrant gene
expression patterns.

[0212] Lipin deficiency attenuates obesity in a genetic model of obesity (Figures
5A, 5B, and 5C). When the fld mutation is transferred into a genetically obese
background, such as the leptin deficient ob/ob mouse strain, resulting lipin- and leptin-
15 double mutant mice exhibit increased food intake equivalent to ob/ob mice, but fail to
become obese. At 14 weeks of age, ob/ob mice reach a weight of 50 grams. In contrast,
mice homozygous for both the ob and fld mutations weigh to only 25 grams;
indistinguishable from wild-type mice.

[0213] Lipin functions at a stage distinct from expansion of the preadipocyte
20 lineage in the maturation of adipocytes. Lipin deficient mice exhibit normal cellularity of
adipose fat pads, but fail to accumulate normal amounts of lipid in cells. Thus, lipin is not
required for expansion of preadipocytes, but for the maturation steps that lead to formation
of normal, lipid-filled adipocytes.

[0214] The data from transgenic mice indicate that lipin Overexpression is
25 associated with increased fat mass. We generated a transgenic mouse strain expressing
lipin specifically in adipose tissue. These mice exhibited a 60% increase in epididymal fat
pad mass on a wild-type genetic background and a 35% increase on an fld strain
background. These results indicate that, whereas lipin deficiency leads to a lack of normal
adipose tissue formation, lipin expression levels above normal can promote excess adipose
30 tissue development. Thus, genetic variation in lipin expression levels can be associated

with either increased or decreased adipose tissue mass, and modulation of lipin levels can be of utility in conditions such as lipodystrophy and obesity.

[0215] Lipin deficiency results in an inherent inability of preadipocytes to differentiate in vitro. Adipogenesis has been examined in fibroblasts isolated from 14 day fld/fld embryos compared to those isolated from wt/fld embryos. When stimulated to differentiate with hormones (insulin, dexamethasone, methylisobutylxanthine) and the PPAR γ ligand rosiglitazone, the fld/fld fibroblasts exhibit reduced lipid accumulation and reduced secretion of lipoprotein lipase activity compared to wt/fld cells. Thus lipin deficiency blocks adipocyte differentiation in a cell autonomous manner.

10 [0216] The lipin protein sequence bears little resemblance to proteins for which function as been previously elucidated. However, there are putative functional motifs that can be identified which we have tested. We find that lipin expression and/or activity can be controlled at several levels, including subcellular localization, protein phosphorylation, and generation of mRNA splice variants that give rise to protein isoforms.

Sub 13
AB
[0217] The N-LIP and acidic lipin protein domains are required for nuclear localization (Figures 6 and 8). Initial studies of lipin localization within the cell revealed that lipin occurs as a predominantly nuclear protein, with a small, but consistent number of cells that exhibit exclusively cytoplasmic localization. Lipin contains a putative nuclear localization signal (NLS) comprised of basic amino acids (KKRRKRRK, SEQ ID NO:___).

20 [0218] Deletion mutations introduced into lipin at other sites indicate that additional lipin protein domains are also required for nuclear localization. The entire region downstream of the NLS, comprising the carboxyl-terminal 75% of the protein and containing the evolutionarily conserved C-LIP domain, is dispensible of nuclear localization. In contrast, deletion of the amino terminal 110 amino acids containing the
25 evolutionarily conserved N-LIP domain abolishes nuclear localization. Deletion of a stretch of ~40 amino acids occurring immediately downstream from the NLS containing a high proportion of acidic amino acid residues also abolishes nuclear localization, whereas deletion of the ~40 amino acid serine-rich domain immediately following the acidic domain does not alter nuclear localization (*see* Figure 8). Thus, lipin contains at least two
30 domains (the N-LIP and acidic domains), in addition to the NLS, that are required for nuclear localization.

[0219] The lipin point mutation occurring in the fld^{2J} mouse strain abolishes nuclear localization. A point mutation in lipin that leads to amino acid substitution at residue 84 (argine substituted for glycine) abolishes nuclear localization. This mutation occurs in the fld^{2J} mouse strain, which shows a similar phenotype to that seen in the fld strain, suggesting that nuclear localization is required for lipin function.

[0220] Lipin gene transcription in adipocytes gives rise to two mRNA splice variants encoding lipin isoforms of potentially different functions. Lipin-1A encodes a predicted 891 amino acid protein, and lipin-1B encodes a predicted 921 amino acid protein resulting from the inclusion of an additional exon encoding 33 amino acids. The lipin-1A and -1B forms may serve different functions in the cell, as the two mRNA variants are present at similar levels in undifferentiated 3T3-L1 preadipocytes, but the expression of lipin-1B increases with differentiation to levels 3-fold higher than lipin-1A after 6 days of differentiation to lipid-containing adipocytes. Furthermore, indirect immunocytochemical studies indicate that lipin-1A is localized primarily to the nucleus, while lipin-1B occurs exclusively in the cytoplasm.

[0221] Within the nucleus, lipin occurs within nuclear speckles, a compartment associated with factors involved in nuclear mRNA metabolism (Figure 7). Lipin that is present in the nucleus exhibits a non-homogenous distribution, suggesting that it may be associated with a subnuclear compartment. It was found that when cells are treated with alpha-amanitin, an inhibitor of transcription, lipin co-localizes with SC-35 (splicing factor, 35 kD) to nuclear speckles. Nuclear speckles typically house proteins having a role in transcriptional regulation or mRNA processing and these results suggest that lipin is involved in mRNA metabolism.

[0222] Lipin is a phosphorylated protein, and the mutant fld^{2J} is aberrantly phosphorylated. The lipin protein sequence contains multiple putative recognition sequences for kinase enzymes. Metabolic labeling of 3T3-L1 and HEK293 cells with ³²P-ATP revealed that both lipin-1A and -1B become phosphorylated, resulting in labeled proteins with retarded electrophoretic mobility. Multiple phosphorylated bands are detected, suggesting that lipin contains multiple phosphorylation sites. A comparison of wild-type lipin to lipin-2J which differs at a single non-phosphorylated amino acid residue (G84R), indicates that lipin-2J is aberrantly phosphorylated as indicated by altered



electrophoretic mobility and fewer phosphorylated protein species. These results suggest that the amino acid change in lipin-2J may affect phosphorylation at distant site, and in light of the exclusion of this form from the nucleus, that phosphorylation may be required for nuclear localization.

- 5 [0223] It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference in their entirety
- 10 for all purposes.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2